

**This dissertation has been  
microfilmed exactly as received 67-17,790**

**KACZYNSKI, Theodore John, 1942-  
BOUNDARY FUNCTIONS.**

**The University of Michigan, Ph.D., 1967  
Mathematics**

**University Microfilms, Inc., Ann Arbor, Michigan**

BOUNDARY FUNCTIONS

by

Theodore John Kaczynski

A dissertation submitted in partial fulfillment  
of the requirements for the degree of  
Doctor of Philosophy in the  
University of Michigan

1967

Doctoral Committee:

Professor Allen L. Shields  
Assistant Professor Peter L. Duren  
Associate Professor Donald J. Livingstone  
Professor Maxwell O. Reade  
Professor Chia-Shun Yih

## BOUNDARY FUNCTIONS

By Theodore John Kaczynski

Let  $H$  denote the set of all points in the Euclidean plane having positive  $y$ -coordinate, and let  $X$  denote the  $x$ -axis. If  $p$  is a point of  $X$ , then by an arc at  $p$  we mean a simple arc  $\gamma$ , having one endpoint at  $p$ , such that  $\gamma - \{p\} \subseteq H$ . Let  $f$  be a function mapping  $H$  into the Riemann sphere. By a boundary function for  $f$  we mean a function  $\varphi$  defined on a set  $E \subseteq X$  such that for each  $p \in E$  there exists an arc  $\gamma$  at  $p$  for which

$$\lim_{\substack{z \rightarrow p \\ z \in \gamma}} f(z) = \varphi(p).$$

The set of curvilinear convergence of  $f$  is the largest set on which a boundary function for  $f$  can be defined; in other words, it is the set of all points  $p \in X$  such that there exists an arc at  $p$  along which  $f$  approaches a limit. A theorem of J. E. McMillan states that if  $f$  is a continuous function mapping  $H$  into the Riemann sphere, then the set of curvilinear convergence of  $f$  is of type  $F_{\delta\delta}$ . In the first of the two chapters of this dissertation we give a more direct proof of this result than McMillan's, and we prove, conversely, that if  $A$  is a set of type  $F_{\delta\delta}$  in  $X$ , then there exists a bounded continuous complex-valued function in  $H$  having  $A$  as its set of curvi-

linear convergence. Next, we prove that a boundary function for a continuous function can always be made into a function of Baire class 1 by changing its values on a countable set of points. Conversely, we show that if  $\varphi$  is a function mapping a set  $E \subseteq X$  into the Riemann sphere, and if  $\varphi$  can be made into a function of Baire class 1 by changing its values on a countable set, then there exists a continuous function in  $H$  having  $\varphi$  as a boundary function. (This is a slight generalisation of a theorem of Bagemihl and Piranian.) In the second chapter we prove that a boundary function for a function of Baire class  $\xi \geq 1$  in  $H$  is of Baire class at most  $\xi+1$ . It follows from this that a boundary function for a Borel-measurable function is always Borel-measurable, but we show that a boundary function for a Lebesgue-measurable function need not be Lebesgue-measurable. The dissertation concludes with a list of problems remaining to be solved.

TABLE OF CONTENTS

	Page
LIST OF ILLUSTRATIONS.....	iii
INTRODUCTION.....	1
Chapter	
I.    BOUNDARY FUNCTIONS FOR CONTINUOUS FUNCTIONS.....	15
II.   BOUNDARY FUNCTIONS FOR DISCONTINUOUS FUNCTIONS.....	57
SOME UNSOLVED PROBLEMS.....	73
REFERENCES.....	75

LIST OF ILLUSTRATIONS

Figure	Page
1. ....	27
2. Trap ( $J, \epsilon, \theta$ ).....	30
3. Tri ( $J, \theta$ ).....	31
4. $S(x_0, \epsilon, \theta)$ .....	32
5. ....	62

## INTRODUCTION

### 1. Preliminary Remarks

Let  $H$  denote the upper half-plane, and let  $X$  denote its frontier, the  $x$ -axis. If  $x \in X$ , then by an arc at  $x$  we mean a simple arc  $\gamma$  with one endpoint at  $x$  such that  $\gamma - \{x\} \subseteq H$ . Suppose that  $f$  is a function mapping  $H$  into some metric space  $Y$ . If  $E$  is any subset of  $X$ , we will say that a function  $\varphi: E \rightarrow Y$  is a boundary function for  $f$  if, and only if, for each  $x \in E$  there exists an arc  $\gamma$  at  $x$  such that

$$\lim_{\substack{z \rightarrow x \\ z \in \gamma}} f(z) = \varphi(x) .$$

The study of boundary functions in this degree of generality was initiated by Bagemihl and Piranian [2]. A function defined in  $H$  may have more than one boundary function defined on a given set  $E \subseteq X$ , but it follows from a famous theorem of Bagemihl [1] that any two such boundary functions differ on at most a countable set of points.

If  $f$  is defined in  $H$ , then the set of curvilinear convergence of  $f$  is the set of all points  $x \in X$  such that there exists some arc at  $x$  along which  $f$  approaches a limit. Evidently, this is the largest set on which a boundary function for  $f$  can be defined.

J. E. McMillan [10] discovered that the set of curvilinear convergence of a continuous function is always of type  $F_{\sigma\delta}$ , and in this paper we show that every set of type  $F_{\sigma\delta}$  in  $X$  is the set of curvilinear

convergence of some continuous function. Next, we show that if  $\varphi$  is a function defined on a subset  $E$  of  $X$ , then  $\varphi$  is a boundary function for some continuous function if and only if  $\varphi$  can be made into a function of the first Baire class by changing its values on at most a countable set of points. (This solves a problem of Bagemihl and Piranian [2, Problem 1].) We then consider functions that are not assumed to be continuous, and we prove that a boundary function for a function of Baire class  $\xi \geq 1$  is of Baire class at most  $\xi + 1$  (thus proving another conjecture of Bagemihl and Piranian [2]). It follows from this that a boundary function for a Borel-measurable function is always Borel-measurable, and in the last section we show that a boundary function for a Lebesgue-measurable function need not be Lebesgue-measurable.

Most of the results appearing here have already been published ([8] and [9]). At the time I published these papers I did not expect to have to make use of this material for a dissertation.

## 2. Notation

$R$  will denote the field of real numbers.

$R^n$  will denote  $n$ -dimensional Euclidean space.

Points in  $R^n$  will be written in the form  $\langle x_1, \dots, x_n \rangle$  rather than  $(x_1, \dots, x_n)$  (to avoid confusion with open intervals of real numbers in the case  $n = 2$ ).

If  $v \in R^n$ , then  $|v|$  denotes the length of the vector  $v$ .

$S^2$  denotes  $\{v \in R^3: |v| = 1\}$ .  $S^2$  will be referred to as the

Riemann sphere.



Let

$$H = \{ \langle x, y \rangle \in \mathbb{R}^2 : y > 0 \}$$

$$H_n = \{ \langle x, y \rangle \in \mathbb{R}^2 : \frac{1}{n} > y > 0 \}$$

$$X = \{ \langle x, 0 \rangle : x \in \mathbb{R} \}$$

$$X_n = \{ \langle x, \frac{1}{n} \rangle : x \in \mathbb{R} \} .$$

We consider  $X$  as being identical with  $\mathbb{R}$ . Thus, for example,  $\langle x, 0 \rangle \leq \langle x', 0 \rangle$  means  $x \leq x'$ , and for  $p, q \in X$ , the notations  $[p, q]$ ,  $[p, q)$ , etc. refer to the obvious intervals on  $X$ .

If  $E$  is a subset of a topological space, then  $\bar{E}$  denotes the closure of  $E$ ,  $E^*$  denotes the interior of  $E$ , and  $E'$  denotes the complement of  $E$ . Of course, if  $E$  is a subset of  $X$ , then  $E^*$  means the interior of  $E$  relative to  $X$ , not relative to the whole plane. In Section 7, we often denote two line segments by  $s$  and  $s'$ . Since the prime notation is never used for complementation in that section, there is no danger of confusing  $s'$  with the complement of  $s$ .

If  $f$  is a function defined in a subset of  $\mathbb{R}^2$ , then  $f(x, y)$  means  $f(\langle x, y \rangle)$ . Thus we write  $f(z)$  for  $z \in \mathbb{R}^2$  and  $f(x, y)$  for  $x, y \in \mathbb{R}$  interchangeably.

### 3. Baire Functions

In this section we review the main facts concerning Borel sets and Baire functions, and we prove some results that will be needed later.

If  $C$  is any family of sets, let  $C_\delta$  be the family of all sets that can be written as a countable intersection of members of  $C$ , and

let  $C_\sigma$  be the family of all sets that can be written as a countable union of members of  $C$ .

Suppose  $M$  is a metrizable topological space. Let  $P^1(M)$  be the family of all open subsets of  $M$  and let  $Q^1(M)$  be the family of all closed subsets of  $M$ . If  $\xi$  is an ordinal number greater than 1, let

$$P^\xi(M) = \left( \bigcup_{\eta < \xi} Q^\eta(M) \right)_\sigma$$

$$Q^\xi(M) = \left( \bigcup_{\eta < \xi} P^\eta(M) \right)_\delta$$

For any  $\xi$ ,  $E \in Q^\xi(M) \iff E' \in P^\xi(M)$ .

For any subset  $L$  of  $M$ ,  $E \in P^\xi(L)$  (respectively  $Q^\xi(L)$ ) if and only if there exists a set  $D \in P^\xi(M)$  (respectively  $Q^\xi(M)$ ) such that  $E = D \cap L$ .

$P^\xi(M)$  and  $Q^\xi(M)$  are closed under finite unions and finite intersections.  $P^\xi(M)$  is closed under countable unions and  $Q^\xi(M)$  is closed under countable intersections.

If  $\eta < \xi$ , then  $P^\eta(M) \cup Q^\eta(M) \subseteq P^\xi(M) \cap Q^\xi(M)$ .

Let  $F_\sigma(M)$  be the class of all  $F_\sigma$  sets of  $M$ , and let  $G_\delta(M)$  be the class of all  $G_\delta$  sets of  $M$ .

$$P^2(M) = F_\sigma(M) \text{ and } Q^2(M) = G_\delta(M).$$

Let  $Y$  be a metric space. For any family  $C$  of subsets of  $M$  we will say that a function  $f : M \rightarrow Y$  is of class  $(C)$  if and only if  $f^{-1}(U) \in C$  for every open set  $U \subseteq Y$ .

The following definition of the Baire classes is somewhat different from the classical definition, but it seems more convenient

for our purposes. A function  $f : M \rightarrow Y$  is said to be of Baire class  $0(M, Y)$  if and only if it is continuous. If  $\xi$  is an ordinal number greater than or equal to 1, then  $f$  is said to be of Baire class  $\xi(M, Y)$  if and only if there exists a sequence of functions  $\{f_n\}_{n=1}^{\infty}$  mapping  $M$  into  $Y$ ,  $f_n$  being of Baire class  $\eta_n(M, Y)$  for some  $\eta_n < \xi$ , such that  $f_n \rightarrow f$  pointwise.

If  $f : M \rightarrow Y$  is of Baire class  $\xi(M, Y)$  and if  $L$  is a subset of  $M$ , then  $f|_L$  is of Baire class  $\xi(L, Y)$ .

If  $K$  is a metric space, if  $g : K \rightarrow M$  is continuous, and if  $f : M \rightarrow Y$  is of Baire class  $\xi(M, Y)$ , then the composite function  $f \circ g$  is of Baire class  $\xi(K, Y)$ .

If  $Y$  is separable and if  $f : M \rightarrow Y$  is of Baire class  $\xi(M, Y)$ , then  $f$  is of class  $(P^{\xi+1}(M))$  [4, page 294].

If  $Y$  is separable and arcwise connected, if  $\xi \neq 1$ , and if  $f : M \rightarrow Y$  is of class  $(P^{\xi+1}(M))$ , then  $f$  is of Baire class  $\xi(M, Y)$  [4].

For any  $\xi$ , if  $f : M \rightarrow R$  is of class  $(P^{\xi+1}(M))$ , then  $f$  is of Baire class  $\xi(M, R)$  [6].

If  $L \in Q^{\xi+1}(M)$  and  $f : L \rightarrow R$  is of Baire class  $\xi(L, R)$ , then  $f$  can be extended to a function  $\bar{f} : M \rightarrow R$  of Baire class  $\xi(M, R)$  [6].

We say that a function  $f : M \rightarrow R$  is Borel measurable if, and only if, for every open set  $U \subseteq R$ ,  $f^{-1}(U)$  is a member of the  $\sigma$ -ring of subsets of  $M$  generated by the open sets.

If  $f : M \rightarrow R$  is of some Baire class  $\xi(M, R)$ , then  $f$  is Borel-measurable, and, conversely, if  $f : M \rightarrow R$  is Borel-measurable, then  $f$  is of Baire class  $\xi(M, R)$  for some countable ordinal number  $\xi$  [7, page 294].

The proofs of Lemmas 1 through 6 are based on standard techniques in the study of Baire functions.

Lemma 1. Let  $M$  be a metric space, and let  $E$  and  $F$  be two  $F_\sigma$  sets in  $M$ . Then there exist two disjoint  $F_\sigma$  sets  $A$  and  $B \subseteq M$  such that

$$E - F \subseteq A \quad \text{and} \quad F - E \subseteq B.$$

Proof. Let  $E = \bigcup_{n=1}^{\infty} E_n$  and  $F = \bigcup_{n=1}^{\infty} F_n$ , where  $E_n$  and  $F_n$  are closed. Then

$$E_n, F_n \in F_\sigma(M) \cap G_\delta(M).$$

It is easy to check that  $F_\sigma(M) \cap G_\delta(M)$  is an algebra (i.e., is closed under complementation, finite unions, and finite intersections). We inductively define a sequence of pairs of sets  $(A_n, B_n)$  as follows. Let

$$A_1 = E_1, \quad B_1 = F_1 \cap A_1'.$$

For  $n > 1$ , let

$$A_n = E_n \cap \bigcap_{j=1}^{n-1} B_j', \quad B_n = F_n \cap \bigcap_{j=1}^n A_j'.$$

By induction,  $A_n, B_n \in F_\sigma(M) \cap G_\delta(M)$ . Let

$$A = \bigcup_{n=1}^{\infty} A_n, \quad B = \bigcup_{n=1}^{\infty} B_n.$$

Then  $A$  and  $B$  are  $F_\sigma$  sets. Notice that

$$\bigcup_{j=1}^{n-1} B_j \subseteq F \quad \text{and} \quad \bigcup_{j=1}^n A_j \subseteq E,$$

from which it follows that

$$A_n = E_n \cap \left( \bigcup_{j=1}^{n-1} B_j \right)' \supseteq E_n \cap F',$$

and

$$B_n = F_n \cap \left( \bigcup_{j=1}^n A_j \right)' \supseteq F_n \cap E'.$$

Therefore

$$A \supseteq \bigcup_{n=1}^{\infty} (E_n \cap F') = E - F$$

and

$$B \supseteq \bigcup_{n=1}^{\infty} (F_n \cap E') = F - E.$$

It only remains to show that  $A \cap B = \phi$ . Suppose  $x \in A \cap B$ . Choose  $\ell, m$  with  $x \in A_\ell$  and  $x \in B_m$ . If  $\ell > m$ , then  $\ell > 1$ , so that

$$A_\ell = E_\ell \cap \bigcap_{j=1}^{\ell-1} B_j' \subseteq B_m'.$$

Hence  $x \in B_m'$  -- a contradiction. On the other hand, if  $\ell \leq m$ , then

$$B_m = F_m \cap \bigcap_{j=1}^m A_j' \subseteq A_\ell'.$$

so that  $x \in A_\ell'$  -- another contradiction. We conclude that  $A \cap B = \phi$ . ■

If  $E$  is a subset of a space  $M$ , we let  $\chi_E$  denote the characteristic function of  $E$ .

Lemma 2. Let  $L$  be a subset of a metric space  $M$ , and suppose that  $E \in F_\sigma(L) \cap G_\delta(L)$ . Then there exists a sequence  $\{f_n\}_{n=1}^{\infty}$  of continuous real-valued functions on  $M$  such that  $f_n \rightarrow \chi_E$  pointwise on  $L$ .

Proof. Both  $E$  and  $L - E$  are in  $F_\sigma(L)$ , so there exist sets  $E_1,$

$F_1 \in F_\sigma(M)$  such that

$$E = E_1 \cap L \quad \text{and} \quad L - E = F_1 \cap L.$$

By Lemma 1, there exist  $A, B \in F_\sigma(M)$  such that  $A \cap B = \phi$  and

$E_1 - F_1 \subseteq A, F_1 - E_1 \subseteq B$ . We have

$$E = A \cap L \quad \text{and} \quad L - E = B \cap L.$$

Write  $A = \bigcup_{n=1}^{\infty} A_n$ ,  $B = \bigcup_{n=1}^{\infty} B_n$ , where  $A_n, B_n$  are closed and  $A_n \subseteq A_{n+1}$ ,  $B_n \subseteq B_{n+1}$  for each  $n$ . By Urysohn's Lemma there exists a continuous function  $f_n : M \rightarrow [0,1]$  such that

$$\begin{aligned} f_n(x) &= 1 && \text{when } x \in A_n \\ f_n(x) &= 0 && \text{when } x \in B_n. \end{aligned}$$

$\{f_n\}_{n=1}^{\infty}$  is the desired sequence. ■

Lemma 3. Let  $L$  be a subset of a metric space  $M$ ,  $f : L \rightarrow R$  a function of class  $(F_{\sigma}(L))$  that takes only finitely many different values. Then there exists a sequence  $\{f_n\}_{n=1}^{\infty}$  of continuous real-valued functions on  $M$  such that  $f_n \rightarrow f$  pointwise on  $L$ .

Proof. From Banach's Hilfssatz 3 [4], we see that there exist real numbers  $a_1, \dots, a_q$  and sets

$$E_1, \dots, E_q \in F_{\sigma}(L) \cap G_{\delta}(L)$$

such that

$$f = \sum_{j=1}^q a_j \chi_{E_j}.$$

If we choose for each  $j$  a sequence  $\{f_n^j\}_{n=1}^{\infty}$  of continuous real-valued functions on  $M$  such that  $f_n^j \rightarrow \chi_{E_j}$  pointwise on  $L$ , and if we set

$$f_n = \sum_{j=1}^q a_j f_n^j,$$

then  $\{f_n\}_{n=1}^{\infty}$  is the desired sequence. ■

Lemma 4. Let  $L$  be a metric space,  $f$  a bounded real-valued function on  $L$  of Baire class 1( $L, R$ ). Then there exists a sequence  $\{f_n\}_{n=1}^{\infty}$

of real-valued functions on  $L$  converging uniformly to  $f$ , such that each  $f_n$  is of class  $(F_\sigma(L))$  and takes only finitely many different values.

Proof.  $f$  is of class  $(F_\sigma(L))$  and the range of  $f$  is totally bounded, so an obvious modification of the proof of Banach's Hilfssatz 4 [4] gives the desired result. ■

Lemma 5. Let  $M$  be a metric space,  $L$  a subset of  $M$ ,  $f : L \rightarrow R$  a function of Baire class 1  $(L, R)$ . Then there exists a sequence  $\{f_n\}_{n=1}^\infty$  of continuous real-valued functions on  $M$  such that  $f_n \rightarrow f$  pointwise on  $L$ .

Proof. We first prove the lemma under the assumption that  $f$  is bounded. For any bounded real-valued function  $h$ , let

$$\|h\| = \sup \{ |h(x)| : x \in \text{domain of } h \}.$$

By Lemma 4 we can choose, for each  $n$ , a function  $g_n : L \rightarrow R$  of class  $(F_\sigma(L))$  such that  $g_n$  takes only finitely many different values and  $\|g_n - f\| \leq \frac{1}{2^n}$ . Let

$$h_1 = g_1, \quad h_n = g_n - g_{n-1} \quad \text{for } n > 1.$$

Then, for  $n > 1$ ,

$$\|h_n\| = \|g_n - f + f - g_{n-1}\| \leq \frac{1}{2^n} + \frac{1}{2^{n-1}} < \frac{1}{2^{n-2}}.$$

Each  $h_n$  is of class  $(F_\sigma(L))$  and takes only finitely many different values, so by Lemma 3 we can choose (for each  $n$ ) a sequence  $\{h_n^j\}_{j=1}^\infty$  of continuous functions on  $M$  such that  $h_n^j \rightarrow h_n$  pointwise on  $L$ .

Set

$$\begin{aligned}
k_n^j(x) &= -\|h_n\| & \text{if } h_n^j(x) \leq -\|h_n\| \\
k_n^j(x) &= \|h_n\| & \text{if } h_n^j(x) \geq \|h_n\| \\
k_n^j(x) &= h_n^j(x) & \text{if } -\|h_n\| < h_n^j(x) < \|h_n\|.
\end{aligned}$$

Then  $k_n^j$  is continuous,  $k_n^j \rightarrow_j h_n$  pointwise on  $L$ , and  $\|k_n^j\| \leq \|h_n\| < \frac{1}{2^{n-2}}$ . Therefore, if we set

$$f_j = \sum_{n=1}^{\infty} k_n^j,$$

then the series converges uniformly and  $f_j$  is continuous on  $M$ . We claim that  $f_j \rightarrow f$  pointwise on  $L$ . Take any  $x \in L$  and any  $\varepsilon > 0$ . Choose  $m$  large enough so that  $\frac{1}{2^{m-2}} < \frac{1}{3} \varepsilon$ . For each  $n$ , choose  $j(n)$  so that

$$j \geq j(n) \Rightarrow |k_n^j(x) - h_n(x)| < \frac{1}{2^{n+1}} \frac{\varepsilon}{3}.$$

Let  $i_0 = \max\{j(1), \dots, j(m)\}$ . Then  $j \geq i_0$  implies that

$$\begin{aligned}
|f_j(x) - f(x)| &\leq |f_j(x) - \sum_{n=1}^m k_n^j(x)| + \left| \sum_{n=1}^m k_n^j(x) - \sum_{n=1}^m h_n(x) \right| \\
&\quad + \left| \sum_{n=1}^m h_n(x) - f(x) \right| \\
&\leq \sum_{n=m+1}^{\infty} \|k_n^j\| + \sum_{n=1}^m |k_n^j(x) - h_n(x)| + \|g_m - f\| \\
&\leq \frac{1}{2^{m-2}} + \left( \sum_{n=1}^m \frac{1}{2^{n+1}} \right) \frac{\varepsilon}{3} + \frac{1}{2^m} < 3 \frac{\varepsilon}{3} = \varepsilon.
\end{aligned}$$

Thus  $f_j(x) \rightarrow_j f(x)$  for each  $x \in L$ , and the lemma is proved for bounded  $f$ .

If  $f$  is not bounded, let



$$g(x) = \arctan f(x) \quad (x \in L).$$

Then  $-\frac{\pi}{2} < g(x) < \frac{\pi}{2}$  for every  $x \in L$ , and  $g$  is of Baire class 1( $L, \mathbb{R}$ ), so there exists a sequence  $\{g_n\}_{n=1}^{\infty}$  of continuous functions on  $M$  converging to  $g$  pointwise on  $L$ . Set

$$\begin{aligned} h_n(x) &= -\frac{\pi}{2} + \frac{1}{n} & \text{if } g_n(x) \leq -\frac{\pi}{2} + \frac{1}{n} \\ h_n(x) &= \frac{\pi}{2} - \frac{1}{n} & \text{if } g_n(x) \geq \frac{\pi}{2} - \frac{1}{n} \\ h_n(x) &= g_n(x) & \text{if } -\frac{\pi}{2} + \frac{1}{n} < g_n(x) < \frac{\pi}{2} - \frac{1}{n}. \end{aligned}$$

Then  $h_n$  is continuous on  $M$ ,  $-\frac{\pi}{2} < h_n(x) < \frac{\pi}{2}$ , and  $h_n \rightarrow g$  pointwise on  $L$ . Let  $f_n(x) = \tan h_n(x)$ . Then  $f_n$  is continuous on  $M$  and  $f_n \rightarrow f$  pointwise on  $L$ . ■

Lemma 6. If  $L$  is a subset of a metric space  $M$  and  $f : L \rightarrow \mathbb{R}^m$  is a function, then the following are equivalent.

- (i)  $f$  is of Baire class 1( $L, \mathbb{R}^m$ ).
- (ii)  $f$  is of class  $(F_\sigma(L))$ .
- (iii) There exists a sequence  $\{f_n\}_{n=1}^{\infty}$  of continuous functions mapping  $M$  into  $\mathbb{R}^m$  such that  $f_n \rightarrow f$  pointwise on  $L$ .

This lemma is an easy consequence of Lemma 5.

Definition. Let  $q$  be any point of  $\mathbb{R}^3$  lying inside the bounded open domain determined by  $S^2$ . By the  $q$ -projection of  $\mathbb{R}^3 - \{q\}$  onto  $S^2$  we mean the function  $P_q$  defined as follows. If  $a$  is any point of  $\mathbb{R}^3 - \{q\}$ , let  $\ell$  be the unique ray, having its endpoint at  $q$ , that passes through  $a$ , and let  $P_q(a)$  be the intersection point of  $\ell$  with  $S^2$ .  $P_q$  is a continuous mapping of  $\mathbb{R}^3 - \{q\}$  onto  $S^2$  that fixes every point of  $S^2$ .

Theorem 1. Let  $L$  be an arbitrary subset of  $\mathbb{R}^2$ . Then a function  $f : L \rightarrow S^2$  is of Baire class 1( $L, S^2$ ) if and only if it is of class  $(F_\sigma(L))$ .

Proof. Assume that  $f : L \rightarrow S^2$  is of class  $(F_\sigma(L))$ .  $S^2 \subseteq \mathbb{R}^3$ , so by Lemma 6 there exists a sequence  $\{f_n\}_{n=1}^\infty$  of continuous functions mapping  $\mathbb{R}^2$  into  $\mathbb{R}^3$  such that  $f_n \rightarrow f$  pointwise on  $L$ . Let

$$A_n = f_n^{-1} \left( \{v \in \mathbb{R}^3 : |v| = \frac{1}{2}\} \right)$$

$$B_n = f_n^{-1} \left( \{v \in \mathbb{R}^3 : |v| \leq \frac{1}{2}\} \right)$$

$$C_n = f_n^{-1} \left( \{v \in \mathbb{R}^3 : |v| \geq \frac{1}{2}\} \right).$$

Let  $f_n^0 = f_n|_{A_n}$ . According to [5, Lemma 2.9, page 299],  $f_n^0$  can be extended to a continuous function  $g_n : \mathbb{R}^2 \rightarrow \{v \in \mathbb{R}^3 : |v| = \frac{1}{2}\}$ .

Define  $h_n : \mathbb{R}^2 \rightarrow \mathbb{R}^3 - \{0\}$  by setting

$$h_n(x) = g_n(x) \quad \text{if } x \in B_n$$

$$h_n(x) = f_n(x) \quad \text{if } x \in C_n.$$

Since  $B_n, C_n$  are closed,  $h_n$  is continuous, and it is easy to verify that  $h_n(x) \rightarrow f(x)$  for each  $x \in L$ . Let  $k_n : \mathbb{R}^2 \rightarrow S^2$  be the composite function  $P_0 \circ h_n$ . Then  $k_n$  is continuous, and for each  $x \in L$ ,  $k_n(x) \rightarrow P_0(f(x)) = f(x)$ . Thus  $f$  is of Baire class 1( $L, S^2$ ). ■

Definition. Let  $M$  and  $Y$  be metric spaces. Then a function  $f : M \rightarrow Y$  is said to be of honorary Baire class 2( $M, Y$ ) if and only if there exists a countable set  $N \subseteq M$  and a function  $g : M \rightarrow Y$  of Baire class 1( $M, Y$ ) such that  $f(x) = g(x)$  for every  $x \in M - N$ .

Theorem 2. Let  $L$  be an arbitrary subset of  $\mathbb{R}^2$  and let  $Y$  be either the real line, a finite-dimensional Euclidean space, or  $S^2$ . Then a

function  $f : L \rightarrow Y$  is of honorary Baire class 2( $L, Y$ ) if and only if there exists a countable set  $N \subseteq L$  such that  $f|_{L-N}$  is of class  $(F_\sigma(L - N))$ .

Proof. Suppose that  $f : L \rightarrow Y$  is of honorary Baire class 2( $L, Y$ ). Then there exists  $g : L \rightarrow Y$  of Baire class 1( $L, Y$ ) and a countable set  $N \subseteq L$  such that  $f|_{L-N} = g|_{L-N}$ . But  $g|_{L-N}$  is of class  $(F_\sigma(L - N))$ .

Conversely, suppose that  $f|_{L-N}$  is of class  $(F_\sigma(L - N))$ , where  $N$  is countable. We must show that  $f$  is of honorary Baire class 2( $L, Y$ ). First consider the case where  $Y = \mathbb{R}^m$ . Write

$$f(x) = \langle f_1(x), f_2(x), \dots, f_m(x) \rangle.$$

Then  $f_i|_{L-N}$  is of class  $(F_\sigma(L - N))$  ( $i=1, \dots, m$ ), and it follows that  $f_i|_{L-N}$  is of Baire class 1( $L - N, \mathbb{R}$ ). Since  $L - N \in G_\delta(L)$ , we can extend  $f_i|_{L-N}$  to a function  $g_i : L \rightarrow \mathbb{R}$  of Baire class 1( $L, \mathbb{R}$ ). If we set  $g(x) = \langle g_1(x), \dots, g_m(x) \rangle$ , then  $g$  is of Baire class 1( $L, \mathbb{R}^m$ ) and  $g(x) = f(x)$  for  $x \in L - N$ , so we have the desired result.

Now consider the case where  $Y = S^2$ . Since  $S^2 \subseteq \mathbb{R}^3$ , there exists, as we have just shown, a function  $g : L \rightarrow \mathbb{R}^3$  of Baire class 1( $L, \mathbb{R}^3$ ) such that  $g(x) = f(x)$  for all  $x \in L - N$ . Then  $g(L) - S^2$  is countable, so there exists some point  $q$  in the bounded open domain determined by  $S^2$  such that  $q \notin g(L)$ . Let  $h$  be the composite function  $P_q \circ g$ . Then  $h$  maps  $L$  into  $S^2$ , and for each  $x \in L - N$ ,

$$h(x) = P_q(g(x)) = P_q(f(x)) = f(x).$$

If  $U \subseteq S^2$  is open, then

$$h^{-1}(U) = g^{-1}(P_q^{-1}(U)) \in F_\sigma(L),$$

so  $h$  is of class  $(F_\sigma(L))$ . By Theorem 1,  $h$  is of Baire class 1( $L, S^2$ ), so we have the desired result. ■

## CHAPTER I

### BOUNDARY FUNCTIONS FOR CONTINUOUS FUNCTIONS

If  $r$  is a positive number and if  $y_0$  is a point of a metric space  $Y$  having metric  $\rho$ , then

$S(r, y_0)$  denotes  $\{y \in Y : \rho(y, y_0) < r\}$ .

We will repeatedly make use of Theorem 11.8 on page 119 in [11] without making explicit reference to it. This theorem states that if  $D$  is a Jordan domain in  $R^2$  or in  $R^2 \cup \{\infty\}$ , if  $\gamma$  is the frontier of  $D$ , and if  $\alpha$  is a cross-cut in  $D$  whose endpoints divide  $\gamma$  into arcs  $\gamma_1$  and  $\gamma_2$ , then  $D - \alpha$  has two components, and the frontiers of these components are respectively  $\alpha \cup \gamma_1$  and  $\alpha \cup \gamma_2$ . (The term cross-cut is defined on page 118 in [11].)

#### 4. Domain of the Boundary Function

**Definition.** If  $f$  is a function mapping into a metric space  $Y$ , then the set of curvilinear convergence of  $f$  is defined to be

$\{x \in X : \text{there exists an arc } \gamma \text{ at } x \text{ and there exists } y \in Y$   
such that

$$\lim_{\substack{z \rightarrow x \\ z \in \gamma}} f(z) = y\}.$$

J. E. McMillan [10] proved that for suitable spaces  $Y$ , the set of curvilinear convergence of a continuous function is always of type  $F_{\sigma\delta}$ . We give a more direct proof of this result than McMillan's. (This proof can be modified to give a more general result; see [9].)

An interval of  $X$  will be called nondegenerate if and only if it contains more than one point.

Suppose  $\gamma$  is a cross-cut of  $H$ . If  $V$  is the bounded component of  $H - \gamma$ , let  $L(\gamma) = \bar{V} \cap X$ . Then  $L(\gamma) = [c, d]$ , where  $c$  and  $d$  are the endpoints of  $\gamma$  and  $c < d$ . Suppose  $\Omega$  is a domain contained in  $H$ . Let  $\Gamma$  denote the family of all cross-cuts  $\gamma$  of  $H$  for which  $\gamma \cap H \subseteq \Omega$ , and let

$$I(\Omega) = \bigcup_{\gamma \in \Gamma} L(\gamma)^*.$$

Let  $\text{acc}(\Omega)$  denote the set of all points on  $X$  that are accessible by arcs in  $\Omega$ .

Lemma 7. Assume that  $\text{acc}(\Omega)$  is nonempty. Let  $a$  be the infimum of  $\text{acc}(\Omega)$  and let  $b$  be the supremum of  $\text{acc}(\Omega)$ . Then

$$I(\Omega) = (a, b).$$

Proof. Suppose  $x \in I(\Omega)$ . Let  $\gamma$  be a cross-cut of  $H$  such that  $\gamma \cap H \subseteq \Omega$  and  $x \in L(\gamma)^*$ .  $L(\gamma) = [c, d]$ , where  $c$  and  $d$  are the endpoints of  $\gamma$  and  $c < d$ . It is evident that  $c$  and  $d$  are in  $\text{acc}(\Omega)$ , so  $a \leq c < x < d \leq b$ , and  $x \in (a, b)$ . Conversely, suppose  $x' \in (a, b)$ . Then there exist points  $c', d' \in \text{acc}(\Omega)$  with  $c' < x' < d'$ . Since  $\Omega$  is arcwise connected, it is easy to show that there exists a cross-cut  $\gamma'$  of  $H$ , with  $\gamma' \cap H \subseteq \Omega$ , having  $c', d'$  as its endpoints. But then  $x' \in (c', d') = L(\gamma')^*$ , so  $x' \in I(\Omega)$ . ■

Lemma 8. If  $\Omega_1$  and  $\Omega_2$  are domains contained in  $H$ , and if

$$(1) \quad I(\Omega_1) \cap \overline{\text{acc}(\Omega_1)} \quad \text{and} \quad I(\Omega_2) \cap \overline{\text{acc}(\Omega_2)}$$

are not disjoint, then  $\Omega_1$  and  $\Omega_2$  are not disjoint.

Proof. We assume that  $\Omega_1$  and  $\Omega_2$  are disjoint and derive a contradiction. Let  $a$  be a point in both of the two sets (1). Let  $\gamma_i$  be a cross-cut of  $H$ , with  $\gamma_i \cap H \subseteq \Omega_i$ , such that  $a \in L(\gamma_i)^*$  ( $i = 1, 2$ ). Let  $U_i$  and  $V_i$  be the components of  $H - \gamma_i$ , where  $V_i$  is the bounded component. Observe that  $\gamma_1 \cap H$  and  $\gamma_2 \cap H$  are disjoint.

Suppose  $\gamma_1 \cap H \subseteq V_2$  and  $\gamma_2 \cap H \subseteq V_1$ . Then, since  $\gamma_1 \cap H \subseteq \bar{U}_1$ ,  $U_1$  has a point in common with  $V_2$ . But, since  $U_1$  is unbounded,  $U_1$  cannot be contained in  $V_2$ , so  $U_1$  must have a point in common with  $\gamma_2 \cap H$ . This contradicts the assumption that  $\gamma_2 \cap H \subseteq V_1$ , so we conclude that either  $\gamma_1 \cap H \not\subseteq V_2$  or  $\gamma_2 \cap H \not\subseteq V_1$ . Hence, either  $\gamma_1 \cap H \subseteq U_2$  or  $\gamma_2 \cap H \subseteq U_1$ . By symmetry, we may assume that  $\gamma_2 \cap H \subseteq U_1$ .

$\Omega_2$  does not meet  $\gamma_1$ , and  $\Omega_2$  does meet  $U_1$  (because  $\gamma_2 \cap H \subseteq U_1 \cap \Omega_2$ ), so  $\Omega_2 \subseteq U_1$ . Since  $a \in \overline{\text{acc}(\Omega_2)}$ , there exists a point  $b \in L(\gamma_1)^*$  such that  $b \in \text{acc}(\Omega_2)$ . But then  $b \in \bar{\Omega}_2 \subseteq \bar{U}_1$ , and this is impossible because the frontier of  $U_1$  is disjoint from  $L(\gamma_1)^*$ . ■

Theorem 3 (J. E. McMillan). Let  $Y$  be a complete separable metric space and let  $f : H \rightarrow Y$  be a continuous function. Then the set of curvilinear convergence of  $f$  is of type  $F_{\alpha \delta}$ .

Proof. Let  $\{p_k\}_{k=1}^{\infty}$  be a countable dense subset of  $Y$ . Let  $\{Q(n, m)\}_{m=1}^{\infty}$  be a counting of all sets of the form

$$\{ \langle x, y \rangle : 0 < y < \frac{1}{n} \text{ and } r < t < r + \frac{1}{n} \}$$

where  $r$  is a rational number. Let  $\{U(n, m, k, \ell)\}_{\ell=1}^{\infty}$  be a counting (with repetitions allowed) of the components of

$$f^{-1}\left(S\left(\frac{1}{2^n}, p_k\right)\right) \cap Q(n, m).$$

(We consider  $\phi$  to be a component of  $\phi$ .) Let

$$A(n, m, k, \ell) = \text{acc}[U(n, m, k, \ell)].$$

Set

$$B = \bigcap_{n=1}^{\infty} \bigcup_{m=1}^{\infty} \bigcup_{k=1}^{\infty} \bigcup_{\ell=1}^{\infty} I(U(n, m, k, \ell)) \cap \overline{A(n, m, k, \ell)}.$$

Since  $I(U(n, m, k, \ell))$  is open in  $X$  it is of type  $F_{\sigma}$ . It follows that  $B$  is of type  $F_{\sigma\delta}$ . Let  $C$  denote the set of curvilinear convergence of  $f$ . I claim that  $B \subseteq C$ . Take any  $b \in B$ . For each  $n$ , choose  $m[n]$ ,  $k[n]$ ,  $\ell[n]$  with

$$(2) \quad b \in I(U(n, m[n], k[n], \ell[n])) \cap \overline{A(n, m[n], k[n], \ell[n])} \\ (n = 1, 2, 3, \dots).$$

For convenience, set  $U_n = U(n, m[n], k[n], \ell[n])$ . By (2) and Lemma 8,  $U_n$  and  $U_{n+1}$  have some point  $z_n$  in common. For each  $n$ , we can choose an arc  $\gamma_n \subseteq U_{n+1}$  with one endpoint at  $z_n$  and the other at  $z_{n+1}$ . Then  $\gamma_n \subseteq Q(n+1, m[n+1])$ . Also,

$$b \in \overline{A(n+1, m[n+1], k[n+1], \ell[n+1])} \subseteq \overline{U_{n+1}} \subseteq \overline{Q(n+1, m[n+1])},$$

and therefore each point of  $\gamma_n$  has distance less than  $\frac{2}{n+1}$  from  $b$ .

$\frac{2}{n+1} \rightarrow 0$  as  $n \rightarrow \infty$ ; hence, if we set  $\gamma = \{b\} \cup \bigcup_{n=1}^{\infty} \gamma_n$ , then  $\gamma$  is an arc with one endpoint at  $b$ .

Since  $U_n$  and  $U_{n+1}$  have a point in common,

$$f^{-1}\left(S\left(\frac{1}{2^n}, p_{k[n]}\right)\right) \text{ and } f^{-1}\left(S\left(\frac{1}{2^{n+1}}, p_{k[n+1]}\right)\right)$$

have a common point, and hence



$$S\left(\frac{1}{2^n}, P_{k[n]}\right) \quad \text{and} \quad S\left(\frac{1}{2^{n+1}}, P_{k[n+1]}\right)$$

have a common point. Therefore, if  $\rho$  is the metric on  $Y$ , then

$$\rho(P_{k[n]}, P_{k[n+1]}) \leq \frac{1}{2^n} + \frac{1}{2^{n+1}} < \frac{1}{2^{n-1}},$$

and therefore

$$\rho(P_{k[n]}, P_{k[n+r]}) \leq \sum_{i=1}^r \rho(P_{k[n+i-1]}, P_{k[n+i]}) < \sum_{i=1}^r \frac{1}{2^{n+i-2}} < \frac{1}{2^{n-2}}.$$

Thus  $\{P_{k[n]}\}$  is a Cauchy sequence and must converge to some point  $p \in Y$ . Since

$$\gamma_n \subseteq U_{n+1} \subseteq f^{-1}\left(S\left(\frac{1}{2^{n+1}}, P_{k[n+1]}\right)\right) \quad \text{and}$$

$$P_{k[n]} \xrightarrow[n]{} p,$$

$\lim_{z \rightarrow b} f(z) = p$ . It is possible that  $\gamma$  is not a simple arc, but according to [12] we can replace  $\gamma$  by a simple arc  $\gamma' \subseteq \gamma$ . Thus  $b \in C$ , and we have shown that  $B \subseteq C$ .

Suppose  $c \in C$ . Let  $\gamma_0$  be an arc at  $c$  such that  $f$  approaches a limit  $p'$  along  $\gamma_0$ . Take any  $n$ . Choose  $k$  with  $p' \in S\left(\frac{1}{2^n}, P_k\right)$ . Choose  $m$  so that  $c$  is in the interior of  $\overline{Q(n, m)} \cap X$ . Then  $\gamma_0$  has a subarc  $\gamma_0'$ , with one endpoint at  $c$ , such that

$$\gamma_0' - \{c\} \subseteq Q(n, m) \cap f^{-1}\left(S\left(\frac{1}{2^n}, P_k\right)\right).$$

Hence, for some  $\ell$ ,  $c \in \text{acc}[U(n, m, k, \ell)] = A(n, m, k, \ell)$ . This shows that

$$C \subseteq \bigcap_{n=1}^{\infty} \bigcup_{m=1}^{\infty} \bigcup_{k=1}^{\infty} \bigcup_{\ell=1}^{\infty} A(n, m, k, \ell).$$

It is easy to deduce from Lemma 7 that the set

$$A(n, m, k, \ell) - I(U(n, m, k, \ell)) = \frac{\dots\dots\dots}{A(n, m, k, \ell) - [I(U(n, m, k, \ell)) \cap A(n, m, k, \ell)]}$$

contains at most two points. It follows by a routine argument that

$$\bigcap_n \bigcup_{m,k,\ell} A(n, m, k, \ell) - \bigcap_n \bigcup_{m,k,\ell} [I(U(n, m, k, \ell)) \cap \overline{A(n, m, k, \ell)}]$$

is countable. Since

$$\bigcap_n \bigcup_{m,k,\ell} [I(U(n, m, k, \ell)) \cap \overline{A(n, m, k, \ell)}] = B \subseteq C$$

$$\bigcap_n \bigcup_{m,k,\ell} A(n, m, k, \ell),$$

$C - B$  is countable, and therefore  $C$  is of type  $F_{\sigma\delta}$ . ■

Next we will show that the foregoing theorem is as strong as possible, in this sense: if  $A$  is any set of type  $F_{\sigma\delta}$  contained in  $X$ , then there exists a bounded continuous complex-valued function  $f$  defined in  $H$  such that  $A$  is the set of curvilinear convergence of  $f$ . The proof is unfortunately quite long.

Definition. Let  $E_1$  and  $E_2$  be two sets on the real line. A point  $p$  on the real line will be called a splitting point for  $E_1$  and  $E_2$  if either

$$x_1 \leq p \text{ for all } x_1 \in E_1 \quad \text{and} \quad p \leq x_2 \text{ for all } x_2 \in E_2$$

$$\text{or} \quad x_2 \leq p \text{ for all } x_2 \in E_2 \quad \text{and} \quad p \leq x_1 \text{ for all } x_1 \in E_1.$$

We will say that two sets  $E_1$  and  $E_2$  split, or that  $E_1$  splits with  $E_2$ , if and only if there exists a splitting point for  $E_1$  and  $E_2$ .

Lemma 9. Let  $E$  be an  $F_\sigma$  set in  $R$ . Then there is a sequence  $\{E_n\}_{n=1}^\infty$  of sets such that

(i)  $E_n$  is bounded and closed

(ii) if  $n \neq m$ , then either  $E_n$  and  $E_m$  are disjoint or  $E_n$  and

$E_m$  split

$$(iii) E = \bigcup_{n=1}^{\infty} E_n.$$

Proof. We can write  $E = \bigcup_{n=1}^{\infty} A_n$  where  $A_n$  is closed,  $A_n \subseteq A_{n+1}$  for all  $n$ , and  $A_1 = \phi$ .

Observe that if  $I$  is any open interval, then there exists a countable family  $\{J_n\}_{n=1}^{\infty}$  of bounded closed intervals such that  $n \neq m \Rightarrow J_n$  and  $J_m$  split, and  $I = \bigcup_{n=1}^{\infty} J_n$ . Since any open set of real numbers is a countable disjoint union of open intervals, it follows that for any open  $U$  there exists a countable family  $\{I_n\}_{n=1}^{\infty}$  of bounded closed intervals such that  $n \neq m \Rightarrow I_n$  and  $I_m$  split, and  $U = \bigcup_{n=1}^{\infty} I_n$ .

For each  $n$ , let  $\{I_j^n\}_{j=1}^{\infty}$  be a family of bounded closed intervals such that  $j \neq k \Rightarrow I_j^n$  and  $I_k^n$  split, and  $A_n' = \bigcup_{j=1}^{\infty} I_j^n$ . Let

$$\mathcal{F} = \{A_1\} \cup \{I_j^n \cap A_{n+1} : n = 1, 2, \dots; j = 1, 2, \dots\}.$$

Then  $\mathcal{F}$  is a countable family of bounded closed sets, and

$$\begin{aligned} E &= \bigcup_{n=1}^{\infty} A_n = A_1 \cup \bigcup_{n=1}^{\infty} (A_{n+1} \cap A_n') \\ &= A_1 \cup \bigcup_{n=1}^{\infty} (A_{n+1} \cap \bigcup_{j=1}^{\infty} I_j^n) = A_1 \cup \bigcup_{n=1}^{\infty} \bigcup_{j=1}^{\infty} (A_{n+1} \cap I_j^n) \\ &= \bigcup \mathcal{F}. \end{aligned}$$

Let  $F_1$  and  $F_2$  be any two distinct members of  $\mathcal{F}$ . If either  $F_1$  or  $F_2$  is  $A_1 = \phi$ , then  $F_1$  and  $F_2$  are automatically disjoint. If neither  $F_1$  nor  $F_2$  is  $A_1$ , then we can write

$$F_1 = I_{j(1)}^{n(1)} \cap A_{n(1)+1} \text{ and } F_2 = I_{j(2)}^{n(2)} \cap A_{n(2)+1}.$$

If  $n(1) < n(2)$ , then  $n(1) + 1 \leq n(2)$ , so

$F_2 = I_{j(2)}^{n(2)} \cap A_{n(2)+1} \subseteq A_{n(2)}' \subseteq A_{n(1)+1}'$ , and therefore  $F_1$  and  $F_2$  are disjoint. A similar argument shows that if  $n(2) < n(1)$ , then  $F_1$  and  $F_2$  are disjoint. Thus, if  $F_1$  and  $F_2$  are not disjoint, then  $n(1) = n(2)$  and we have

$$F_1 = I_{j(1)}^n \cap A_{n+1} \text{ and } F_2 = I_{j(2)}^n \cap A_{n+1},$$

where  $n = n(1) = n(2)$ . But then  $j(1) \neq j(2)$ , so  $I_{j(1)}^n$  and  $I_{j(2)}^n$  split, and therefore  $F_1$  and  $F_2$  split. So we have shown that any two distinct members of  $\mathcal{F}$  either split or are disjoint.

If  $\mathcal{F}$  has infinitely many distinct members, let  $E_1, E_2, E_3, \dots$  be a counting of  $\mathcal{F}$ . If  $\mathcal{F}$  has only finitely many distinct members, let  $E_1, \dots, E_m$  be the members of  $\mathcal{F}$  and let  $E_k = \phi$  for  $k > m$ . In either case,  $\{E_n\}_{n=1}^{\infty}$  is the desired sequence. ■

If  $F$  is a closed subset of the real line, then by a complementary interval of  $F$  we mean a component of  $F'$ . (If  $F = \mathbb{R}$ , then  $\phi$  is considered to be a complementary interval of  $F$ .)

Definition. By a special family we mean a family  $\mathcal{F}$  of subsets of  $\mathbb{R}$  such that

- (3)  $\mathcal{F}$  is nonempty
- (4) each member of  $\mathcal{F}$  is bounded and closed
- (5) there exists a sequence  $\{F_n\}_{n=1}^{\infty}$  of members of  $\mathcal{F}$  such that every member of  $\mathcal{F}$  is equal to some  $F_n$ , and the following condition is satisfied.

(5a) If  $m > n$ , then either  $F_m$  is contained in one of the complementary

intervals of  $F_n$ , or else  $F_m$  splits with  $F_n$ .

Lemma 10. If  $E$  is an  $F_\sigma$  set in  $R$ , then there exists a special family  $\mathcal{F}$  such that  $E = \bigcup \mathcal{F}$ .

Proof. By Lemma 9 we can choose a sequence  $\{E_n\}_{n=1}^\infty$  of bounded closed sets such that if  $n \neq m$  then  $E_n$  and  $E_m$  either split or are disjoint, and  $E = \bigcup_{n=1}^\infty E_n$ .

Let  $n_1 = 1$  and let  $F_1 = E_1$ . Now suppose that  $n_1, n_2, \dots, n_s$  are chosen and  $F_1, F_2, \dots, F_{n_s}$  are chosen so that

$$(i) \quad 1 = n_1 < n_2 < \dots < n_s$$

$$(ii) \quad F_i \text{ is closed and bounded } (i = 1, \dots, n_s)$$

(iii) if  $n_s \geq r > t \geq 1$ , then either  $F_r$  is contained in one of the complementary intervals of  $F_t$ , or else  $F_r$  splits with  $F_t$

(iv) if  $1 \leq i \leq n_s$ , then there exists  $j \in \{1, \dots, s\}$  such that  $F_i \subseteq E_j$

$$(v) \quad \bigcup_{i=1}^{n_s} F_i = \bigcup_{i=1}^s E_i.$$

We construct  $F_{n_s+1}, \dots, F_{n_{s+1}}$  as follows. Let  $\mathcal{I}$  be the family of complementary intervals of the bounded closed set

$$\bigcup_{i=1}^{n_s} F_i = \bigcup_{i=1}^s E_i.$$

We assert that  $E_{s+1}$  meets at most finitely many members of  $\mathcal{I}$ . If this assertion is false, then there exists an infinite sequence  $\{I_n\}_{n=1}^\infty$  of members of  $\mathcal{I}$  such that  $n \neq m$  implies  $I_n \cap I_m = \emptyset$ , and there exists (for each  $m$ ) a point  $x_m \in I_m \cap E_{s+1}$ .  $\{x_m\}_{m=1}^\infty$  is a bounded sequence, and  $n \neq m$  implies that  $x_n \neq x_m$ . From this it follows that  $\{x_m\}_{m=1}^\infty$  has either a strictly increasing or a strictly decreasing convergent

subsequence. We will assume that  $\{x_{m(k)}\}_{k=1}^{\infty}$  is a strictly increasing convergent subsequence; the reasoning is similar in the case of a strictly decreasing convergent subsequence. Say  $I_{m(k)} = (a_k, b_k)$ . Then  $a_k < x_{m(k)} < b_k$ , so since  $x_{m(k)} < x_{m(k+1)} < b_{k+1}$  and  $x_{m(k)} \notin I_{m(k+1)}$  we must have  $x_{m(k)} \leq a_{k+1} < x_{m(k+1)}$ . Therefore, if we let

$$\bar{x} = \lim_{k \rightarrow \infty} x_{m(k)},$$

then  $\bar{x} = \lim_{k \rightarrow \infty} a_k$  also. Moreover, for  $k \geq 2$ ,  $a_k$  is a finite real number, so that  $a_k \in \bigcup_{i=1}^s E_i$ . Therefore there exists  $u \in \{1, \dots, s\}$  such that  $a_k \in E_u$  for infinitely many values of  $k$ . Consequently  $\bar{x} \in E_u$ . But since  $x_{m(k)} \in E_{s+1}$ ,  $\bar{x} \in E_{s+1}$  also. But then  $\bar{x} \in E_u \cap E_{s+1}$ , so that  $E_u$  and  $E_{s+1}$  must split and  $\bar{x}$  must be a splitting point for  $E_u$  and  $E_{s+1}$ . Since infinitely many  $a_k$  lie in  $E_u$ ,  $E_u$  contains points that are less than  $\bar{x}$ ; and  $E_{s+1}$  also contains points less than  $\bar{x}$ ; therefore  $E_u$  and  $E_{s+1}$  cannot split, and we have a contradiction. This proves the assertion. **Let**

$$\mathcal{J} = \{\phi\} \cup \{\bar{I} \cap E_{s+1} : I \in \mathcal{I} \text{ and } I \cap E_{s+1} \neq \phi\}.$$

Let  $n_{s+1}$  equal  $n_s$  plus the number of members of  $\mathcal{J}$ . Let  $F_{n_{s+1}}, \dots, F_{n_{s+1}}$  be all the members of  $\mathcal{J}$ . We must show that conditions (i) through (v) are still satisfied when  $s$  is replaced by  $s+1$ . Conditions (i), (ii), and (iv) are obvious. The verification of (iii) is divided into three parts. Suppose  $n_{s+1} \geq r > t \geq 1$ .

Case I. Assume that  $n_s \geq r > t \geq 1$ . In this case we already know that either  $F_r$  is contained in one of the complementary intervals of  $F_t$  or else  $F_r$  splits with  $F_t$ .

Case II. Assume that  $n_{s+1} \geq r > n_s \geq t \geq 1$ . There exists  $v \in \{1, \dots, s\}$  such that  $F_t \subseteq E_v$ . Either  $E_v$  and  $E_{s+1}$  are disjoint or they split.

Case IIa. Assume  $E_v$  and  $E_{s+1}$  are disjoint. Either  $F_r = \phi$  (in which case  $F_r$  is certainly contained in a complementary interval of  $F_t$ ) or else  $F_r \neq \phi$  and  $F_r = \bar{I} \cap E_{s+1}$  for some  $I \in \mathcal{Q}$ . Let  $J$  be the smallest closed interval containing  $F_r$ . Then  $J \subseteq \bar{I}$  and  $J^* \subseteq I \subseteq (\bigcup_{i=1}^s E_i)'$ , so that  $J^*$  does not meet  $E_v$ . The endpoints of  $J$  lie in  $F_r \subseteq E_{s+1}$ , so neither endpoint of  $J$  lies in  $E_v$ . So  $J$  does not meet  $E_v$  and therefore  $J$  does not meet  $F_t$ ; from which it follows that  $F_r$  is contained in a complementary interval of  $F_t$ .

Case IIb. Assume that  $E_v$  and  $E_{s+1}$  split. Since  $F_t \subseteq E_v$  and  $F_r \subseteq E_{s+1}$  it follows that  $F_t$  and  $F_r$  split.

Case III. Assume that  $n_{s+1} \geq r > t > n_s$ . If either  $F_r$  or  $F_t$  is  $\phi$ , it is clear that  $F_r$  is contained in a complementary interval of  $F_t$ . Otherwise, there exist  $I_1, I_2 \in \mathcal{Q}$  such that  $I_1 \wedge I_2 = \phi$  and

$$F_r = \bar{I}_1 \cap E_{s+1} \quad \text{and} \quad F_t = \bar{I}_2 \cap E_{s+1}.$$

Since  $\bar{I}_1$  and  $\bar{I}_2$  evidently split,  $F_r$  and  $F_t$  must split.

Thus condition (iii) is verified.

As for (v), it is clear that

$$E_{s+1} - \bigcup_{i=1}^s E_i \subseteq \bigcup_{j=n_s+1}^{n_{s+1}} F_j \subseteq E_{s+1},$$

so that

$$\begin{aligned} \bigcup_{i=1}^{s+1} E_i &= \left( \bigcup_{i=1}^s E_i \right) \cup \left( E_{s+1} - \bigcup_{i=1}^s E_i \right) \\ &\subseteq \left( \bigcup_{j=1}^{n_s} F_j \right) \cup \left( \bigcup_{j=n_s+1}^{n_{s+1}} F_j \right) \subseteq \left( \bigcup_{i=1}^s E_i \right) \cup E_{s+1} = \bigcup_{i=1}^{s+1} E_i. \end{aligned}$$

$$\text{Hence } \bigcup_{i=1}^{s+1} E_i = \left( \bigcup_{j=1}^{n_s} F_j \right) \cup \left( \bigcup_{j=n_s+1}^{n_{s+1}} F_j \right) = \bigcup_{j=1}^{n_{s+1}} F_j.$$

Thus we have shown that we can construct sequences  $\{n_j\}_{j=1}^{\infty}$ ,  $\{F_k\}_{k=1}^{\infty}$  in such a way that conditions (i) through (v) are satisfied for every value of  $s$ . If we set  $\mathcal{F} = \{F_k : k = 1, 2, \dots\}$ , it is easy to verify that  $\mathcal{F}$  is a special family and that  $E = \bigcup \mathcal{F}$ . ■

Definition. If  $\mathcal{F}_1$  and  $\mathcal{F}_2$  are two families of sets, let

$$\mathcal{F}_1 \wedge \mathcal{F}_2 = \{F_1 \cap F_2 : F_1 \in \mathcal{F}_1 \text{ and } F_2 \in \mathcal{F}_2\}.$$

Lemma 11. If  $\mathcal{F}$  and  $\tilde{\mathcal{F}}$  are two special families, then  $\mathcal{F} \wedge \tilde{\mathcal{F}}$  is a special family.

Proof. Conditions (3) and (4) in the definition of a special family are clearly satisfied, so we just have to verify (5).

Arrange all pairs of positive integers in a sequence according to the scheme shown in Figure 1. Let  $(a(k), b(k))$  be the  $k$ th term of the sequence<sup>1</sup> ( $k = 1, 2, \dots$ ). Observe that  $k < \ell$  if and

<sup>1</sup>The reader may find it amusing to derive the following formulas for  $(a(k), b(k))$ . For real  $t$ , let  $[[t]]$  denote the largest integer that is strictly less than  $t$ . Then

$$\begin{aligned} a(k) &= \frac{1}{2} \left( \left[ \left[ \frac{\sqrt{8k+1} + 1}{2} \right] \right]^2 + \left[ \left[ \frac{\sqrt{8k+1} + 1}{2} \right] \right] \right) - k + 1 \\ &= \frac{1}{8} \left( \left[ \left[ \sqrt{8k+1} \right] \right] + \frac{5}{2} - \frac{1}{2}(-1)^{\left[ \left[ \sqrt{8k+1} \right] \right]} \right) \left( \left[ \left[ \sqrt{8k+1} \right] \right] + \frac{1}{2} - \frac{1}{2}(-1)^{\left[ \left[ \sqrt{8k+1} \right] \right]} \right) - k + 1 \\ &= \begin{cases} \frac{1}{8} \left( \left[ \left[ \sqrt{8k+1} \right] \right] + 3 \right) \left( \left[ \left[ \sqrt{8k+1} \right] \right] + 1 \right) - k + 1 & \text{if } \left[ \left[ \sqrt{8k+1} \right] \right] \text{ is odd} \\ \frac{1}{8} \left( \left[ \left[ \sqrt{8k+1} \right] \right] + 2 \right) \left[ \left[ \sqrt{8k+1} \right] \right] - k + 1 & \text{if } \left[ \left[ \sqrt{8k+1} \right] \right] \text{ is even,} \end{cases} \end{aligned}$$

and






Figure 1.

only if either  $a(k) + b(k) < a(\ell) + b(\ell)$  or else  $a(k) + b(k) = a(\ell) + b(\ell)$  and  $b(k) < b(\ell)$ . Thus  $k < \ell$  implies that either  $a(k) < a(\ell)$  or  $b(k) < b(\ell)$ .

Let  $\{F_n\}_{n=1}^{\infty}$  be a sequence of elements of  $\mathcal{F}$  such that every member of  $\mathcal{F}$  is equal to some  $F_n$  and such that condition (5a) in the definition of a special family is satisfied. Let  $\{\tilde{F}_n\}_{n=1}^{\infty}$  be a similar sequence for  $\tilde{\mathcal{F}}$ . Set

$$\hat{F}_k = F_{a(k)} \cap \tilde{F}_{b(k)}.$$

Then  $\{\hat{F}_k\}_{k=1}^{\infty}$  is a sequence in  $\mathcal{F} \wedge \tilde{\mathcal{F}}$  such that every member of  $\mathcal{F} \wedge \tilde{\mathcal{F}}$  is equal to some  $\hat{F}_k$ . We must show that condition (5a) is satisfied. Suppose that  $\ell > k$ . Two cases occur.

Case I.  $a(k) < a(\ell)$ .

Note that  $\hat{F}_k \subseteq F_{a(k)}$  and  $\hat{F}_\ell \subseteq F_{a(\ell)}$ . Either  $F_{a(\ell)}$  is contained in one of the complementary intervals of  $F_{a(k)}$  (in which case  $\hat{F}_\ell$  is contained in a complementary interval of  $\hat{F}_k$ ), or else  $F_{a(\ell)}$  and  $F_{a(k)}$  split (in which case  $\hat{F}_\ell$  and  $\hat{F}_k$  split).

Case II.  $b(k) < b(\ell)$ .

In this case a similar argument shows that either  $\hat{F}_\ell$  is contained in

$$\begin{aligned} b(k) &= \frac{1}{2} \left( \left[ \left[ \frac{\sqrt{8k+1} + 1}{2} \right] \right] - \left[ \left[ \frac{\sqrt{8k+1} + 1}{2} \right] \right]^2 \right) + k \\ &= \frac{-1}{8} \left( \left[ \left[ \sqrt{8k+1} \right] \right] + \frac{1}{2} - \frac{1}{2}(-1)^{\left[ \left[ \sqrt{8k+1} \right] \right]} \right) \left( \left[ \left[ \sqrt{8k+1} \right] \right] - \frac{3}{2} - \frac{1}{2}(-1)^{\left[ \left[ \sqrt{8k+1} \right] \right]} \right) + k \\ &= \begin{cases} \frac{-1}{8} \left( \left[ \left[ \sqrt{8k+1} \right] \right] + 1 \right) \left( \left[ \left[ \sqrt{8k+1} \right] \right] - 1 \right) + k & \text{if } \left[ \left[ \sqrt{8k+1} \right] \right] \text{ is odd} \\ \frac{-1}{8} \left[ \left[ \sqrt{8k+1} \right] \right] \left( \left[ \left[ \sqrt{8k+1} \right] \right] - 2 \right) + k & \text{if } \left[ \left[ \sqrt{8k+1} \right] \right] \text{ is even.} \end{cases} \end{aligned}$$

a complementary interval of  $\hat{F}_k$  or  $\hat{F}_l$  and  $\hat{F}_k$  split. Thus condition (5a) is satisfied, and  $\mathcal{F} \wedge \tilde{\mathcal{F}}$  is a special family. ■

Lemma 12. Let  $E_1, E_2$  be two  $F_\sigma$  sets in  $R$  such that  $E_1 \subseteq E_2$ , and suppose that  $\mathcal{F}_1$  and  $\mathcal{F}_2$  are special families such that  $E_1 = \bigcup \mathcal{F}_1$  and  $E_2 = \bigcup \mathcal{F}_2$ . Then  $E_1 = \bigcup (\mathcal{F}_1 \wedge \mathcal{F}_2)$ .

The proof is obvious.

Next we introduce some notation.

Let  $J$  be a nonempty interval on  $X$  with endpoints  $a, b$  ( $a \leq b$ ).

By  $\text{Trap}(J, \varepsilon, \theta)$  (where  $\theta \in (0, \frac{\pi}{2})$  and  $\varepsilon > 0$ ) we mean the interior of the trapezoid shown in Figure 2. That is,

$$\text{Trap}(J, \varepsilon, \theta) = \{ \langle x, y \rangle : 0 < y < \varepsilon, a + y \text{ctn } \theta < x < b - y \text{ctn } \theta \}.$$

For  $\theta \in (0, \frac{\pi}{2})$ , let  $\text{Tri}(J, \theta)$  be the closed triangular area shown in Figure 3. That is,

$$\text{Tri}(J, \theta) = \{ \langle x, y \rangle : y \geq 0 \text{ and } a + y \text{ctn } \theta \leq x \leq b - y \text{ctn } \theta \}.$$

If  $x_0 \in X$ ,  $\varepsilon > 0$ , and  $\theta \in (0, \frac{\pi}{2})$ , let  $S(x_0, \varepsilon, \theta)$  denote the open Stolz angle shown in Figure 4. That is,

$$S(x_0, \varepsilon, \theta) = \{ \langle x, y \rangle : 0 < y < \varepsilon, \\ x_0 + y \text{ctn } (\pi - \theta) < x < x_0 + y \text{ctn } \theta \}.$$

If  $K$  is a closed set on a real line, let  $J(K)$  be the smallest closed interval containing  $K$ . If  $K$  is bounded, closed, and nonempty,  $\varepsilon > 0$ , and  $0 < \beta < \alpha < \frac{\pi}{2}$ , then we define

$$B(K, \varepsilon, \alpha, \beta) = \text{Trap}(J(K), \varepsilon, \alpha) - \bigcup_{I \in \mathcal{I}} \text{Tri}(I, \beta),$$

where  $\mathcal{I}$  denotes the set of complementary intervals of  $K$ .




Figure 2.--Trap( $J, \epsilon, \theta$ )




Figure 3.--Tri(J,  $\theta$ )




Figure 4.-- $S(x_0, \epsilon, \theta)$

We state without proof the following readily verifiable facts.

(6)  $B(K, \varepsilon, \alpha, \beta)$  is an open subset of  $H$ .

(7)  $S(e, \varepsilon, \theta)$  is an open subset of  $H$ .

(8) If  $K_1$  and  $K_2$  split, then for any  $\varepsilon_1, \varepsilon_2, \alpha, \beta$ ,

$$B(K_1, \varepsilon_1, \alpha, \beta) \quad \text{and} \quad B(K_2, \varepsilon_2, \alpha, \beta)$$

are disjoint.

(9) Suppose that  $K_1 \subseteq K$ ,  $\varepsilon > \varepsilon_1 > 0$ , and  $0 < \beta < \beta_1 < \alpha_1 < \alpha < \frac{\pi}{2}$ .

Then

$$\overline{B(K_1, \varepsilon_1, \alpha_1, \beta_1)} \cap H \subseteq B(K, \varepsilon, \alpha, \beta).$$

(10) Suppose  $K_1$  is contained in one of the complementary intervals of  $K$ , and suppose  $\varepsilon, \alpha, \beta$  are given. Then there exists  $\delta > 0$  such that for every  $\eta \leq \delta$ ,

$$B(K, \varepsilon, \alpha, \beta) \quad \text{and} \quad B(K_1, \eta, \alpha, \beta)$$

are disjoint.

(11) Suppose that  $\alpha < \theta < \frac{\pi}{2}$  and  $x_0 \notin J(K)^*$ . Then, for any  $\varepsilon, \varepsilon_1, \beta$ ,

$$B(K, \varepsilon, \alpha, \beta) \quad \text{and} \quad S(x_0, \varepsilon_1, \theta)$$

are disjoint.

(12) Suppose that  $x_0 \in K \cap J(K)^*$  and  $\beta < \alpha < \theta < \frac{\pi}{2}$ . Let  $\varepsilon$  be given.

Then there exists  $\delta > 0$  such that for every  $\eta \leq \delta$ ,

$$\overline{S(x_0, \eta, \theta)} \cap H \subseteq B(K, \varepsilon, \alpha, \beta).$$

(13) Suppose that  $\varepsilon < \varepsilon'$  and  $\theta' < \theta$ . Then

$$\overline{S(x_0, \varepsilon, \theta)} \cap H \subseteq S(x_0, \varepsilon', \theta').$$

- (14) Suppose  $x_0 \notin K$  and  $\varepsilon, \alpha, \beta, \theta$  are given. Then there exists  $\delta > 0$  such that for every  $\eta \leq \delta$ ,

$$S(x_0, \eta, \theta) \quad \text{and} \quad B(K, \varepsilon, \alpha, \beta)$$

are disjoint.

- (15) If  $x_0 \neq x_1$  and  $\varepsilon, \theta$  are given, then there exists  $\delta > 0$  such that for every  $\eta \leq \delta$ ,

$$S(x_0, \varepsilon, \theta) \quad \text{and} \quad S(x_1, \eta, \theta)$$

are disjoint.

(16)  $\overline{B(K, \varepsilon, \alpha, \beta)} \cap X \subseteq K.$

(17)  $\overline{S(x_0, \varepsilon, \theta)} \cap X = \{x_0\}.$

Definition. If  $\mathcal{F}$  is a special family, let  $\mathcal{F}^2$  be the set of all members of  $\mathcal{F}$  that have two or more points.

Definition. Let  $\mathcal{F}$  be a special family, let  $E$  be the set of all endpoints of intervals  $J(F)$  where  $F \in \mathcal{F}$ ,  $F \neq \phi$ , and suppose that  $0 < \beta < \alpha < \theta < \frac{\pi}{2}$ . By a pair of special  $\alpha, \beta, \theta$  functions for  $\mathcal{F}$  we mean a pair  $(\varepsilon, \delta)$ , where  $\varepsilon$  and  $\delta$  are positive real-valued functions, the domain of  $\varepsilon$  is  $E$ , the domain of  $\delta$  is  $\mathcal{F}^2$ , and

- (18) for each  $\eta > 0$ , there exist at most finitely many  $F \in \mathcal{F}^2$  such that  $\delta(F) \geq \eta$ ;
- (19) for each  $\eta > 0$ , there exist at most finitely many  $e \in E$  such that  $\varepsilon(e) \geq \eta$ ;
- (20) if  $e, e' \in E$  and  $e \neq e'$ , then

$$S(e, \varepsilon(e), \theta) \quad \text{and} \quad S(e', \varepsilon(e'), \theta)$$



are disjoint;

(21) if  $F, K \in \mathcal{F}^2$  and  $F \neq K$ , then

$$B(F, \delta(F), \alpha, \beta) \quad \text{and} \quad B(K, \delta(K), \alpha, \beta)$$

are disjoint;

(22) if  $e \in E$  and  $F \in \mathcal{F}^2$ , then

$$S(e, \varepsilon(e), \theta) \quad \text{and} \quad B(F, \delta(F), \alpha, \beta)$$

are disjoint.

Lemma 13. Let  $\mathcal{F}$  be a special family and suppose that  $0 < \beta < \alpha < \theta < \frac{\pi}{2}$ . Then there exists a pair of special  $\alpha, \beta, \theta$  functions for  $\mathcal{F}$ .

Proof. Let  $\{F_n\}_{n=1}^{\infty}$  be a sequence of members of  $\mathcal{F}$  of the type referred to in condition (5) in the definition of a special family. Let

$$\mathcal{F}^2(n) = \{F \in \mathcal{F}^2 : F = F_k \text{ for some } k \leq n\}$$

$E$  = set of all endpoints of intervals  $J(F)$  for

$$F \in \mathcal{F}, F \neq \phi$$

$E(n) = \{e \in E : e \text{ is an endpoint of } J(F_k) \text{ for}$

some  $k \leq n$  for which  $F_k \neq \phi\}$ .

If  $J(F_1)$  has one endpoint  $e$ , set  $\varepsilon(e) = 1$ . If  $J(F_1)$  has two endpoints  $e_1, e_2$ , then by (15) we can choose  $\varepsilon(e_1) \leq 1$  and  $\varepsilon(e_2) \leq 1$  so that  $S(e_1, \varepsilon(e_1), \theta)$  and  $S(e_2, \varepsilon(e_2), \theta)$  are disjoint. If  $F_1 \in \mathcal{F}^2$ , set  $\delta(F_1) = 1$ . In this case,  $J(F_1)$  has two endpoints  $e_1$  and  $e_2$  and (by (11))  $B(F_1, \delta(F_1), \alpha, \beta)$ ,  $S(e_1, \varepsilon(e_1), \theta)$ , and  $S(e_2, \varepsilon(e_2), \theta)$  are all disjoint.

Now suppose that  $\varepsilon(e)$  and  $\delta(F)$  have been defined for all

$e \in E(n)$  and all  $F \in \mathcal{F}^2(n)$  in such a way that

(i) if  $e, e' \in E(n)$  and  $e \neq e'$ , then  $S(e, \varepsilon(e), \theta)$  and  $S(e', \varepsilon(e'), \theta)$  are disjoint;

(ii) if  $F, K \in \mathcal{F}^2(n)$  and  $F \neq K$ , then  $B(F, \delta(F), \alpha, \beta)$  and  $B(K, \delta(K), \alpha, \beta)$  are disjoint;

(iii) if  $e \in E(n)$  and  $F \in \mathcal{F}^2(n)$ , then  $S(e, \varepsilon(e), \theta)$  and  $B(F, \delta(F), \alpha, \beta)$  are disjoint;

(iv) if  $e \in E(n)$  and  $k \leq n$  is the least integer for which  $e \in E(k)$ , then  $\varepsilon(e) \leq \frac{1}{k}$ ;

(v) if  $F \in \mathcal{F}^2(n)$  and  $k \leq n$  is the least integer for which  $F \in \mathcal{F}^2(k)$ , then  $\delta(F) \leq \frac{1}{k}$ .

We must extend the definitions of  $\varepsilon$  and  $\delta$  to  $E(n+1)$  and  $\mathcal{F}^2(n+1)$  in such a way that conditions (i) through (v) are still satisfied when  $n$  is replaced by  $n+1$ .

Case I. If  $F_{n+1} = \phi$  or if  $F_{n+1} = F_k$  for some  $k \leq n$ , then  $\mathcal{F}^2(n+1) = \mathcal{F}^2(n)$  and  $E(n+1) = E(n)$ , so that nothing is required to be done.

Case II. If  $F_{n+1}$  consists of a single point  $e$  and if  $e \in F_k$  for some  $k \leq n$ , then (since  $F_{n+1}$  and  $F_k$  must split in this case)  $e$  is an endpoint of  $J(F_k)$ , so that again  $\mathcal{F}^2(n+1) = \mathcal{F}^2(n)$  and  $E(n+1) = E(n)$ , and nothing is required to be done.

Case III. Suppose that  $F_{n+1}$  consists of a single point  $e_0$  and that for each  $k \leq n$ ,  $e_0 \notin F_k$ . By (14), (15), and the fact that  $E(n)$  and  $\mathcal{F}^2(n)$  are finite, we can choose  $\varepsilon(e_0) \in (0, \frac{1}{n+1})$  so that  $S(e_0, \varepsilon(e_0), \theta)$  is disjoint from  $S(e, \varepsilon(e), \theta)$  and from  $B(F, \delta(F), \alpha, \beta)$  for each  $e \in E(n)$  and each  $F \in \mathcal{F}(n)$ . The construction is then finished for

$E(n+1)$  and  $F(n+1)$ .

Case IV. Suppose that  $F_{n+1}$  contains at least two points and that, for each  $k \leq n$ ,  $F_k \not\perp F_{n+1}$ . For each  $k \leq n$ , either  $F_{n+1}$  splits with  $F_k$ , or else  $F_{n+1}$  is contained in a complementary interval of  $F_k$ . Since  $\mathcal{F}^2(n)$  is finite, (8) and (10) show that we can choose  $\delta(F_{n+1}) \in (0, \frac{1}{n+1})$  so that  $B(F_{n+1}, \delta(F_{n+1}), \alpha, \beta)$  is disjoint from  $B(F, \delta(F), \alpha, \beta)$  for each  $F \in \mathcal{F}^2(n)$ .

Say  $e \in E(n)$ . Then  $e$  is an endpoint of  $J(F_k)$  for some  $k \leq n$ , so (since  $F_{n+1}$  either splits with  $F_k$  or is contained in a complementary interval of  $F_k$ )  $e \notin J(F_{n+1})^*$ . By (11),  $B(F_{n+1}, \delta(F_{n+1}), \alpha, \beta)$  and  $S(e, \varepsilon(e), \theta)$  are disjoint.

Let  $e_0, e_0'$  be the endpoints of  $J(F_{n+1})$ .

Case IVa.  $e_0, e_0' \in E(n)$ .

In this case the construction is already finished.

Case IVb.  $e_0 \in E(n)$  and  $e_0' \notin E(n)$ .

If  $e_0' \in F_k$  for some  $k \leq n$ , then  $F_{n+1}$  splits with  $F_k$ , so that  $e_0'$  must be an endpoint of  $J(F_k)$  -- which contradicts the assumption that  $e_0' \notin E(n)$ . Hence, for each  $k \leq n$ ,  $e_0' \notin F_k$ . By (14), (15), and the fact that  $E(n)$  and  $\mathcal{F}^2(n)$  are finite, we can choose  $\varepsilon(e_0') \in (0, \frac{1}{n+1})$  so that  $S(e_0', \varepsilon(e_0'), \theta)$  is disjoint from  $S(e, \varepsilon(e), \theta)$  and from  $B(F, \delta(F), \alpha, \beta)$  for each  $e \in E(n)$  and each  $F \in \mathcal{F}^2(n)$ . By (11),  $S(e_0', \varepsilon(e_0'), \theta)$  and  $B(F_{n+1}, \delta(F_{n+1}), \alpha, \beta)$  are disjoint. Thus the construction is finished for  $E(n+1)$  and  $\mathcal{F}^2(n+1)$ .

Case IVc.  $e_0 \notin E(n)$  and  $e_0' \in E(n)$ .

This case is essentially the same as Case IVb.

Case IVd.  $e_0 \notin E(n)$  and  $e_0' \notin E(n)$ .

If  $e_0 \in F_k$  for some  $k \leq n$ , then  $F_{n+1}$  splits with  $F_k$ , so  $e_0$  is an endpoint of  $J(F_k)$ ; a contradiction. Thus  $e_0 \notin F_k$  for  $k \leq n$ , and similarly  $e_0' \notin F_k$  for  $k \leq n$ . Therefore, by (14) and (15), we can choose  $\varepsilon(e_0)$  and  $\varepsilon(e_0') \in (0, \frac{1}{n+1})$  so that  $S(e_0, \varepsilon(e_0), \theta)$  and  $S(e_0', \varepsilon(e_0'), \theta)$  are disjoint and each of  $S(e_0, \varepsilon(e_0), \theta)$  and  $S(e_0', \varepsilon(e_0'), \theta)$  is disjoint from every  $S(e, \varepsilon(e), \theta)$  ( $e \in E(n)$ ) and from every  $B(F, \delta(F), \alpha, \beta)$  ( $F \in \mathcal{F}^2(n)$ ). By (11),  $S(e_0, \varepsilon(e_0), \theta)$  and  $S(e_0', \varepsilon(e_0'), \theta)$  are each disjoint from  $B(F_{n+1}, \delta(F_{n+1}), \alpha, \beta)$ , so the construction is finished for  $E(n+1)$  and  $\mathcal{F}^2(n+1)$ .

We have shown that we can inductively define  $\varepsilon(e)$  for every  $e \in E$  and  $\delta(F)$  for every  $F \in \mathcal{F}^2$  in such a way that (i) through (v) are satisfied for every value of  $n$ . Conditions (20), (21) and (22) in the definition of a pair of  $\alpha, \beta, \theta$  special functions are thus automatically satisfied by  $(\varepsilon, \delta)$ . We must verify that (18) and (19) are also satisfied.

Suppose (19) is false. Then there exists  $\eta > 0$  and there exists an infinite sequence  $\{e_k\}_{k=1}^{\infty}$  of distinct members of  $E$  such that  $\varepsilon(e_k) \geq \eta$  for every  $k$ . Let  $m(k)$  be the least integer for which  $e_k$  is an endpoint of  $J(F_{m(k)})$ . Each  $J(F_m)$  has at most two endpoints, so, since the  $e_k$  are all distinct, there exists (for given  $m$ ) at most two values of  $k$  for which  $m(k) = m$ . Therefore there exist infinitely many distinct integers among  $m(1), m(2), m(3), \dots$ . Consequently there exists  $j$  with  $\frac{1}{m(j)} < \eta$ . But, by (iv),  $\varepsilon(e_j) \leq \frac{1}{m(j)} < \eta$ , a contradiction. So (19) must be true. A similar argument shows that (18) is true. ■

Lemma 14. Let  $\mathcal{F}$  be a special family,  $0 < \beta < \alpha < \theta < \frac{\pi}{2}$ , and let  $E$  be the set of all endpoints of intervals  $J(F)$  for  $F \in \mathcal{F}$ . Suppose  $(\varepsilon, \delta)$  is a pair of special  $\alpha, \beta, \theta$  functions for  $\mathcal{F}$ . If  $\varepsilon_1, \delta_1$  are two real-valued functions having domains  $E$  and  $\mathcal{F}^2$  respectively, and if

$$\begin{aligned} 0 < \varepsilon_1(e) &\leq \varepsilon(e) && \text{for all } e \in E, \text{ and} \\ 0 < \delta_1(F) &\leq \delta(F) && \text{for all } F \in \mathcal{F}^2, \end{aligned}$$

then  $(\varepsilon_1, \delta_1)$  is a pair of special  $\alpha, \beta, \theta$  functions for  $\mathcal{F}$ .

Proof. This follows from the fact that

$$S(x_0, \varepsilon', \theta) \subseteq S(x_0, \varepsilon'', \theta)$$

and

$$B(K, \varepsilon', \alpha, \beta) \subseteq B(K, \varepsilon'', \alpha, \beta)$$

whenever  $\varepsilon' \leq \varepsilon''$ . ■

Theorem 4. Let  $A$  be any set of type  $F_{\sigma\delta}$  in  $X$ . Then there exists a bounded continuous complex-valued function  $f$  defined in  $H$  such that  $A$  is the set of curvilinear convergence of  $f$ .

Proof. We can write  $A = \bigcap_{n=1}^{\infty} A_n$ , where each  $A_n$  is of type  $F_{\sigma}$  and  $A_{n+1} \subseteq A_n$  for every  $n$ . For each  $n$ , let  $\mathcal{F}_n$  be a special family with  $\bigcup \mathcal{F}_n = A_n$ . Let

$$\begin{aligned} \hat{\mathcal{F}}_1 &= \mathcal{F}_1 \\ \hat{\mathcal{F}}_{n+1} &= \hat{\mathcal{F}}_n \wedge \mathcal{F}_{n+1} \quad \text{for } n > 1. \end{aligned}$$

By Lemmas 11 and 12, together with mathematical induction,  $\hat{\mathcal{F}}_n$  is a special family and  $\bigcup \hat{\mathcal{F}}_n = A_n$ . Moreover, every member of  $\hat{\mathcal{F}}_{n+1}$  is a subset of some member of  $\hat{\mathcal{F}}_n$ .

Let  $\{\beta_n\}_{n=1}^{\infty}$  be a strictly ascending sequence in  $(0, \frac{\pi}{8})$  converging to  $\frac{\pi}{8}$ .

Let  $\{\alpha_n\}_{n=1}^{\infty}$  be a strictly descending sequence in  $(\frac{\pi}{8}, \frac{\pi}{4})$  converging to  $\frac{\pi}{8}$ .

Let  $\{\theta_n\}_{n=1}^{\infty}$  be a strictly ascending sequence in  $(\frac{\pi}{4}, \frac{3\pi}{8})$  converging to  $\frac{3\pi}{8}$ .

Let  $E_n$  be the set of all endpoints of intervals  $J(F)$  for  $F \in \hat{\mathcal{F}}_n$ .

Let  $(\varepsilon(1, \cdot), \delta(1, \cdot))$  be any pair of special  $\alpha_1, \beta_1, \theta_1$  functions for  $\hat{\mathcal{F}}_1$ .

Now suppose that for each  $k \leq n$  we have chosen a pair of special  $\alpha_k, \beta_k, \theta_k$  functions  $(\varepsilon(k, \cdot), \delta(k, \cdot))$  for  $\hat{\mathcal{F}}_k$  in such a way that

(i) whenever  $1 \leq k \leq n - 1$ ,  $e \in E_{k+1}$ ,  $F \in \hat{\mathcal{F}}_k^2$ , and  $e \in F \cap J(F)^*$ , then

$$\overline{S(e, \varepsilon(k+1, e), \theta_{k+1})} \cap H \subseteq B(F, \delta(k, F), \alpha_k, \beta_k);$$

(ii) whenever  $1 \leq k \leq n - 1$ ,  $e \in E_{k+1}$ , and  $e \in E_k$ , then

$$\overline{S(e, \varepsilon(k+1, e), \theta_{k+1})} \cap H \subseteq S(e, \varepsilon(k, e), \theta_k);$$

(iii) whenever  $1 \leq k \leq n - 1$ ,  $K \in \hat{\mathcal{F}}_{k+1}^2$ ,  $F \in \hat{\mathcal{F}}_k^2$ , and  $K \subseteq F$ , then

$$\overline{B(K, \delta(k+1, K), \alpha_{k+1}, \beta_{k+1})} \cap H \subseteq B(F, \delta(k, F), \alpha_k, \beta_k).$$

Then we construct  $(\varepsilon(n+1, \cdot), \delta(n+1, \cdot))$  as follows. Let  $(\varepsilon, \delta)$  be any pair of special  $\alpha_{n+1}, \beta_{n+1}, \theta_{n+1}$  functions for  $\hat{\mathcal{F}}_{n+1}$ . If  $e \in E_{n+1} - E_n$ , then for some unique  $F \in \hat{\mathcal{F}}_n^2$ ,  $e \in F \cap J(F)^*$ , so by (12) we can choose  $\xi(e) > 0$  such that  $\eta \leq \xi(e)$  implies

$$\overline{S(e, \eta, \theta_{n+1})} \cap H \subseteq B(F, \delta(n, F), \alpha_n, \beta_n).$$

We set  $\varepsilon(n+1, e) = \min \{\varepsilon(e), \xi(e)\}$ . On the other hand, if  $e \in E_{n+1} \cap E_n$ ,

then we set  $\varepsilon(n+1, e) = \min \{ \varepsilon(e), \frac{1}{2} \varepsilon(n, e) \}$ .

If  $F \in \hat{\mathcal{F}}_{n+1}^2$ , then there exists a unique  $K \in \hat{\mathcal{F}}_n^2$  with  $F \subseteq K$ .

Set

$$\delta(n+1, F) = \min \{ \delta(F), \frac{1}{2} \delta(n, K) \}.$$

By Lemma 14,  $(\varepsilon(n+1, \cdot), \delta(n+1, \cdot))$  is a pair of special  $\alpha_{n+1}$ ,  $\beta_{n+1}$ ,  $\theta_{n+1}$  functions for  $\hat{\mathcal{F}}_{n+1}$ , and by (13) and (9), conditions (i), (ii), and (iii) are still satisfied when  $n$  is replaced by  $n+1$ . Thus we can inductively construct a pair  $(\varepsilon(n, \cdot), \delta(n, \cdot))$  of special  $\alpha_n$ ,  $\beta_n$ ,  $\theta_n$  functions for  $\hat{\mathcal{F}}_n$  in such a way that conditions (i), (ii) and (iii) are satisfied for every  $n$ .

Let

$$U_n = \left[ \bigcup_{e \in E_n} S(e, \varepsilon(n, e), \theta_n) \right] \cup \left[ \bigcup_{F \in \hat{\mathcal{F}}_n^2} B(F, \delta(n, F), \alpha_n, \beta_n) \right].$$

Then  $U_n$  is open. For fixed  $n$ , all the various sets  $S(e, \varepsilon(n, e), \theta_n)$  ( $e \in E_n$ ) and  $B(F, \delta(n, F), \alpha_n, \beta_n)$  ( $F \in \hat{\mathcal{F}}_n^2$ ) are open and pairwise disjoint, so that every component of  $U_n$  is contained in one of the sets  $S(e, \varepsilon(n, e), \theta_n)$  ( $e \in E_n$ ) or  $B(F, \delta(n, F), \alpha_n, \beta_n)$  ( $F \in \hat{\mathcal{F}}_n^2$ ). It therefore follows from (16) and (17) that if  $\Omega$  is any component of  $U_n$ , then

$$(23) \quad \bar{\Omega} \cap X \subseteq A_n.$$

From the fact that  $(\varepsilon(n, \cdot), \delta(n, \cdot))$  is a pair of special  $\alpha_n$ ,  $\beta_n$ ,  $\theta_n$  functions for  $\hat{\mathcal{F}}_n$  together with conditions (18) and (19), it follows that

$$\bar{U}_n \cap H = \left[ \bigcup_{e \in E_n} \overline{S(e, \varepsilon(n, e), \theta_n)} \cap H \right] \cup \left[ \bigcup_{F \in \mathcal{F}_n^2} \overline{B(F, \delta(n, F), \alpha_n, \beta_n)} \cap H \right].$$

Consequently, conditions (i), (ii), (iii), together with the fact that

$$e \in E_{n+1} - E_n \Rightarrow e \in F \cap J(F)^* \text{ for some } F \in \mathcal{F}_n^2,$$

show that  $\bar{U}_{n+1} \cap H \subseteq U_n$  for every  $n$ .

By Urysohn's Lemma, there exists a continuous function  $g_n : H \rightarrow [0, 1]$  such that

$$g_n(z) = 1 \text{ for } z \in H - U_n$$

and

$$g_n(z) = 0 \text{ for } z \in \bar{U}_{n+1} \cap H.$$

Let  $g(z) = \sum_{n=1}^{\infty} \frac{1}{2^n} g_n(z)$ . Then  $0 \leq g(z) \leq 1$ , and the series converges uniformly, so  $g$  is continuous on  $H$ .

If  $z \in H - U_n$ , then  $z \in H - U_m$  for every  $m \geq n$ , so that

$$1 = g_n(z) = g_{n+1}(z) = g_{n+2}(z) = \dots, \text{ and hence}$$

$$(24) \quad g(z) \geq \sum_{m=n}^{\infty} \frac{1}{2^m} = \frac{1}{2^{n-1}} \quad (z \in H - U_n).$$

Also, if  $z \in U_{n+1}$ , then  $z \in U_1, U_2, \dots, U_{n+1}$ , so that

$$0 = g_1(z) = g_2(z) = \dots = g_n(z), \text{ and}$$

$$(25) \quad g(z) \leq \sum_{m=n+1}^{\infty} \frac{1}{2^m} = \frac{1}{2^n} \quad (z \in U_{n+1}).$$

We assert that

$$(26) \quad \text{for each } x_0 \in A, g(z) \rightarrow 0 \text{ as } z \rightarrow x_0 \text{ with } z \in S(x_0, 1, \frac{3\pi}{8}).$$

Take any natural number  $n$ . Since  $x_0 \in A_{n+1} = \bigcup \hat{\mathcal{F}}_{n+1}$ , either  $x_0 \in E_{n+1}$  or else  $x_0 \in F \cap J(F)^*$  for some  $F \in \hat{\mathcal{F}}_{n+1}^2$ . In the first



case, set  $\eta = \varepsilon(n+1, x_0)$ . In the second case, (12) shows that we can choose  $n > 0$  so that

$$S(x_0, \eta, \frac{3\pi}{8}) \subseteq B(F, \delta(n+1, F), \alpha_{n+1}, \beta_{n+1}).$$

Suppose  $\langle x, y \rangle \in S(x_0, 1, \frac{3\pi}{8})$  and  $y < \eta$ . Then, in the first case,  $\langle x, y \rangle \in S(x_0, \eta, \frac{3\pi}{8}) \subseteq S(x_0, \varepsilon(n+1, x_0), \theta_{n+1}) \subseteq U_{n+1}$ , and in the second case,

$$\langle x, y \rangle \in S(x_0, \eta, \frac{3\pi}{8}) \subseteq B(F, \delta(n+1, F), \alpha_{n+1}, \beta_{n+1}) \subseteq U_{n+1}. \text{ So, by (25),}$$

$$\begin{aligned} (\langle x, y \rangle \in S(x_0, 1, \frac{3\pi}{8}) \text{ and } y < \eta) &\implies \langle x, y \rangle \in U_{n+1} \\ &\implies 0 \leq g(x, y) \leq \frac{1}{2^n}. \end{aligned}$$

This proves (26).

Let  $x_0$  be a point in  $X$  and  $\gamma$  any arc at  $x_0$ . Suppose  $g(z) \rightarrow 0$  as  $z \rightarrow x_0$  along  $\gamma$ . Then  $\gamma$  has a subarc  $\gamma'$  with one endpoint at  $x_0$  such that  $\gamma' - \{x_0\} \subseteq g^{-1}((- \frac{1}{2^n}, \frac{1}{2^n})$ . By (24),  $\gamma' - \{x_0\} \subseteq U_n$ . Therefore, by (23),  $x_0 \in A_n$ . Since  $n$  was arbitrary,  $x_0 \in \bigcap_{n=1}^{\infty} A_n = A$ . Thus,

(27) if there exists an arc  $\gamma$  at  $x_0$  such that  $g(z) \rightarrow 0$  as  $z$  approaches  $x_0$  along  $\gamma$ , then  $x_0 \in A$ .

Now define

$$f(x, y) = g(x, y) \sin \frac{1}{y} + i g(x, y) \quad (\langle x, y \rangle \in H).$$

If  $x_0 \in A$ , then, by (26),  $f(z) \rightarrow 0$  as  $z \rightarrow x_0$  with  $z \in S(x_0, 1, \frac{3\pi}{8})$ .

Thus every point of  $A$  is in the set of curvilinear convergence of  $f$ .

Conversely, suppose  $x_0$  is any point of the set of curvilinear convergence of  $f$ . Let  $\gamma$  be an arc at  $x_0$  such that  $f$  approaches the limit  $c + di$  along  $\gamma$ . Then  $g$  approaches the limit  $d$  along  $\gamma$ . If  $d$

is different from zero, then  $g(x, y) \sin \frac{1}{y}$  (the real part of  $f$ ) cannot approach any limit along  $\gamma$  -- a contradiction. Therefore  $g$  approaches the limit 0 along  $\gamma$ , and, by (27),  $x_0 \in A$ . Therefore  $A$  is the set of curvilinear convergence of  $f$ . ■

### 5. Boundary Functions for Continuous Functions

Lemma 15. Let  $E$  be a metric space,  $Y$  a separable metric space.

Suppose that  $\varphi: E \rightarrow Y$  is a function having the following property.

For every open set  $U \subseteq Y$  there exists an  $F_\sigma$  set  $L \subseteq E$  and a countable set  $N \subseteq E$  such that

$$\varphi^{-1}(U) \subseteq L \subseteq \varphi^{-1}(\bar{U}) \cup N.$$

Then there exists a countable set  $M \subseteq E$  such that  $\varphi|_{E-M}$  is of class  $(F_\sigma(E-M))$ .

Proof. Let  $\mathcal{B}$  be a countable base for  $Y$ . For each  $B \in \mathcal{B}$ , let  $L(B) \subseteq E$  be an  $F_\sigma$  set and let  $N(B) \subseteq E$  be a countable set such that

$$\varphi^{-1}(B) \subseteq L(B) \subseteq \varphi^{-1}(\bar{B}) \cup N(B)$$

Let  $M = \bigcup_{B \in \mathcal{B}} N(B)$ . Then  $M$  is countable. Let  $E_0 = E - M$  and let  $\varphi_0 = \varphi|_{E_0}$ . We show that  $\varphi_0$  is of class  $(F_\sigma(E_0))$ .

Let  $W$  be any open subset of  $Y$ . If  $p \in W$ , there exists  $r > 0$  such that  $S(r, p) \subseteq W$ . Choose  $B \in \mathcal{B}$  so that  $p \in B \subseteq S(\frac{1}{2}r, p)$ . Then  $\bar{B} \subseteq S(r, p) \subseteq W$ . It follows that

$$W = \bigcup_{B \in \mathcal{Q}(W)} B = \bigcup_{B \in \mathcal{Q}(W)} \bar{B},$$

where  $\mathcal{Q}(W) = \{B \in \mathcal{B} : \bar{B} \subseteq W\}$ . Therefore

$$\varphi_0^{-1}(W) = E_0 \cap \varphi^{-1}(W) = E_0 \cap \bigcup_{B \in \mathcal{Q}(W)} \varphi^{-1}(B)$$

$$\begin{aligned}
&\subseteq E_0 \cap \bigcup_{B \in \mathcal{Q}(W)} L(B) \\
&\subseteq E_0 \cap \bigcup_{B \in \mathcal{Q}(W)} [\varphi^{-1}(\bar{B}) \cup N(B)] \\
&\subseteq E_0 \cap \bigcup_{B \in \mathcal{Q}(W)} [\varphi^{-1}(\bar{B}) \cup M] \\
&= E_0 \cap \bigcup_{B \in \mathcal{Q}(W)} \varphi^{-1}(\bar{B}) \\
&= E_0 \cap \varphi^{-1}(W) = \varphi_0^{-1}(W).
\end{aligned}$$

Consequently  $\varphi_0^{-1}(W) = E_0 \cap \bigcup_{B \in \mathcal{Q}(W)} L(B)$ , so  $\varphi_0^{-1}(W)$  is of class  $(F_\sigma(E_0))$ . ■

**Theorem 5.** Let  $Y$  be a separable metric space and let  $f : H \rightarrow Y$  be a continuous function. Suppose that  $E \subseteq X$  and that  $\varphi : E \rightarrow Y$  is a boundary function for  $f$ . Then there exists a countable set  $M \subseteq E$  such that  $\varphi|_{E-M}$  is of class  $(F_\sigma(E-M))$ .

**Proof.** Let  $U$  be any open subset of  $Y$ , and let  $W = (\bar{U})'$ . Let

$$E_n = \{x \in X : \text{there exists an arc } \gamma \text{ at } x, \text{ having one endpoint on } X_n, \text{ such that } \gamma - \{x\} \subseteq f^{-1}(U)\}$$

$$K = \{x \in X : \text{there exists an arc } \gamma \text{ at } x \text{ such that } \gamma - \{x\} \subseteq f^{-1}(W)\}.$$

Observe that

$$\varphi^{-1}(U) \subseteq \bigcup_{n=1}^{\infty} E_n,$$

and

$$\varphi^{-1}(W) \subseteq K.$$

For the time being, let  $n$  be a fixed natural number. For each  $x \in K$  we can choose an arc  $\gamma_x$  at  $x$  such that

$$\gamma_x - \{x\} \subseteq H_n \cap f^{-1}(W).$$

Since an arc at  $x$  is by definition a simple arc,  $\gamma_x - \{x\}$  is a connected set and hence must be contained within one nonempty component of  $H_n \cap f^{-1}(W)$ . Let  $U_x$  denote this component (for each  $x \in K$ ).

Let  $T$  be the set of all points of  $K$  that are two-sided limit points of  $\bar{E}_n$ . We claim that if  $x, y \in T$ , then  $x \neq y$  implies  $U_x \cap U_y = \phi$ . If  $U_x \cap U_y \neq \phi$ , then (since  $U_x$  and  $U_y$  are two components of the same set)  $U_x$  and  $U_y$  are equal. Let  $p$  be the endpoint of  $\gamma_x$  lying in  $U_x$  and let  $q$  be the endpoint of  $\gamma_y$  lying in  $U_y = U_x$ . We can join  $p$  and  $q$  by an arc  $\gamma$  lying in  $U_x$ . Putting  $\gamma_x$ ,  $\gamma_y$  and  $\gamma$  together, we obtain an arc  $\alpha$  with one endpoint at  $x$  and the other at  $y$ , such that  $\alpha - \{x, y\} \subseteq U_x$ . According to [12] we can choose a simple arc  $\alpha' \subseteq \alpha$  having one endpoint at  $x$  and the other at  $y$ . Of course,  $\alpha' - \{x, y\} \subseteq U_x \subseteq H_n \cap f^{-1}(W)$ . Let  $I$  be the open interval in  $X$  with endpoints at  $x$  and  $y$ , and let  $J = X - \bar{I}$ . Let  $B$  be the bounded component of  $H - \alpha'$  and let  $A$  be the other component. Since  $X_n$  is unbounded and does not meet  $\alpha'$ ,  $X_n \subseteq A$ .

Because  $x$  is a two-sided limit point of  $\bar{E}_n$ , we can choose a point  $w \in I \cap E_n$ . Let  $\beta$  be an arc at  $w$ , having one endpoint on  $X_n$ , such that  $\beta - \{w\} \subseteq f^{-1}(U)$ . Then  $\beta$  does not meet  $\alpha'$  (because  $\alpha' - \{x, y\} \subseteq f^{-1}(W)$  and  $f^{-1}(W) \cap f^{-1}(U) = \phi$ ), and therefore (since  $\beta - \{w\}$  contains a point of  $X_n \subseteq A$ )  $\beta - \{w\} \subseteq A$ . It follows that  $w \in \bar{A}$ . This, however, is a contradiction, because the frontier of  $A$  (relative to the finite plane) is  $\alpha' \cup J$ . We conclude that, for  $x, y \in T$ ,  $x \neq y$  implies  $U_x \cap U_y = \phi$ .

An open set in the plane has only countably many components, so it follows that  $T$  must be countable. Let  $S$  be the set of all

points of  $\bar{E}_n$  that are not two-sided limit points of  $\bar{E}_n$ . We know that  $S$  is countable, so

$$\begin{aligned} K \cap \bar{E}_n &= [K \cap (\bar{E}_n - S)] \cup [K \cap S] \\ &= T \cup [K \cap S] \end{aligned}$$

is countable.

Let  $N = K \cap \bigcup_{n=1}^{\infty} \bar{E}_n = \bigcup_{n=1}^{\infty} (K \cap \bar{E}_n)$ . Then  $N$  is countable, and, since  $\varphi^{-1}(W) \subseteq K$ ,

$$\begin{aligned} \varphi^{-1}(U) &\subseteq E \cap \bigcup_{n=1}^{\infty} \bar{E}_n \subseteq E \cap \bigcup_{n=1}^{\infty} \bar{E}_n \\ &= (E \cap K \cap \bigcup_{n=1}^{\infty} \bar{E}_n) \cup ((E - K) \cap \bigcup_{n=1}^{\infty} \bar{E}_n) \\ &\subseteq (E \cap N) \cup (E - K) \subseteq (E \cap N) \cup (E - \varphi^{-1}(W)) \\ &= (E \cap N) \cup \varphi^{-1}(\bar{U}). \end{aligned}$$

Thus  $\varphi^{-1}(U) \subseteq E \cap \bigcup_{n=1}^{\infty} \bar{E}_n \subseteq (E \cap N) \cup \varphi^{-1}(\bar{U})$ , and the desired result follows from Lemma 15. ■

Corollary. Let  $Y$  be either the Riemann sphere, the real line, or a finite-dimensional Euclidean space. If  $f : H \rightarrow Y$  is a continuous function, if  $E \subseteq X$ , and if  $\varphi : E \rightarrow Y$  is a boundary function for  $f$ , then  $\varphi$  is of honorary Baire class 2  $(E, Y)$ .

Next we show that the foregoing corollary is as strong as possible in the sense that if  $E$  is any subset of  $X$  and  $\varphi$  is a function of honorary Baire class 2 mapping  $E$  into a suitable space, then there exists a continuous function in  $H$  having  $\varphi$  as a boundary function. A proof of this result -- at least for real- or vector-valued functions -- was outlined by Bagemihl and Piranian [2, Theorem 8], in the case

where  $E = X$ . Although the construction given here is carried out much more explicitly than the construction given by Bagemihl and Piranian, my treatment differs from theirs in only two aspects that are of any significance. First of all, the proof of the theorem for arbitrary subsets  $E$  of  $X$  depends on Lemma 6 of the Introduction. Secondly, Bagemihl and Piranian say in the last line of their proof that there is "no difficulty now in extending  $f$  continuously to the whole of  $D$  in such a manner that  $\phi$  is a boundary function for  $f$ ." While this appears to be all right for real- or vector-valued functions, it is not clear why the extension should be so easy for functions taking values on the Riemann sphere. Theorem 7 of the present paper shows, however, that the result can be obtained for functions taking values on the sphere once it is known for vector-valued functions.

The following miniature closed graph theorem will be a convenience.

Lemma 16. Suppose that  $M$  is a metric space and that  $u : M \rightarrow R$  is a function having the following properties:

(i) if  $\{p_n\}$  is a convergent sequence of points of  $M$ , then  $\{u(p_n)\}$  converges neither to  $+\infty$  nor to  $-\infty$ ;

(ii) if  $\{p_n\} \subseteq M$ ,  $p \in M$ , and  $y \in R$ , and if  $p_n \xrightarrow{n} p$  and  $u(p_n) \xrightarrow{n} y$ , then  $u(p) = y$ .

Then  $u$  is continuous.

Proof. Suppose that  $\{p_n\}$  is a sequence of points in  $M$  converging to a point  $p \in M$ . Using (i) it is easy to show that  $\{u(p_n)\}$  is a bounded sequence. Suppose that  $\{u(p_n)\}$  does not converge to  $u(p)$ . Then there exists a subsequence  $\{u(p_{n(k)})\}$  that converges to a real number  $y \neq u(p)$ . This, however, contradicts (ii). We conclude that

$$u(p_n) \rightarrow u(p). \blacksquare$$

Lemma 17. Let  $h : \mathbb{R} \rightarrow \mathbb{R}$  be a strictly increasing function such that  $h(\mathbb{R})$  is neither bounded above nor bounded below. Then there exists a continuous weakly increasing function  $h^* : \mathbb{R} \rightarrow \mathbb{R}$  such that  $h^*(h(x)) = x$  for every  $x \in \mathbb{R}$ .

Proof. Let  $Z = h(\mathbb{R})$ . Observe that  $h^{-1} : Z \rightarrow \mathbb{R}$  is strictly increasing. For any  $x \in \mathbb{R}$ , the set  $(-\infty, x] \cap Z$  is nonempty. Also,  $h^{-1}((-\infty, x] \cap Z)$  is bounded above, because if we choose  $y \in Z$  with  $x \leq y$ , then  $h^{-1}((-\infty, x] \cap Z)$  is bounded above by  $h^{-1}(y)$ .

We claim that for every  $x \in \mathbb{R}$

$$(27) \quad \sup h^{-1}((-\infty, x] \cap Z) = \sup h^{-1}((-\infty, x) \cap Z).$$

If  $x \notin Z$ , the equation is trivial. Suppose  $x \in Z$ . Then

$$y < h^{-1}(x) \implies (h(y) < x \text{ and } h(y) \in Z),$$

so that  $h((-\infty, h^{-1}(x))) \subseteq (-\infty, x) \cap Z$ . Hence

$$(-\infty, h^{-1}(x)) \subseteq h^{-1}((-\infty, x) \cap Z),$$

so that  $\sup h^{-1}((-\infty, x) \cap Z) \geq h^{-1}(x) = \sup h^{-1}((-\infty, x] \cap Z)$ . The opposite inequality is trivial, so (27) is established.

We also claim that

$$(28) \quad \inf h^{-1}((x, +\infty) \cap Z) = \sup h^{-1}((-\infty, x] \cap Z).$$

Obviously,  $\inf h^{-1}((x, +\infty) \cap Z) \geq \sup h^{-1}((-\infty, x] \cap Z)$ . Take any  $y > \sup h^{-1}((-\infty, x] \cap Z)$ . If  $h(y) \leq x$ , then  $h(y) \in (-\infty, x] \cap Z$ , and so  $y \in h^{-1}((-\infty, x] \cap Z)$ -- a contradiction. Thus  $h(y) > x$  and  $h(y) \in (x, +\infty) \cap Z$ . Therefore  $y \in h^{-1}((x, +\infty) \cap Z)$ , and so  $\inf h^{-1}((x, +\infty) \cap Z) \leq y$ . In view of the choice of  $y$ , this implies

that

$$\inf h^{-1}((x, +\infty) \cap Z) \leq \sup h^{-1}((-\infty, x] \cap Z),$$

and (28) is established.

Define

$$h^*(x) = \sup h^{-1}((-\infty, x] \cap Z).$$

It is clear that  $h^*$  is weakly increasing and that  $h^*(h(x)) = x$  for every real  $x$ . The continuity of  $h^*$  can easily be deduced from the equations

$$\begin{aligned} \sup h^*((-\infty, x)) &= h^*(x) \\ \inf h^*((x, +\infty)) &= h^*(x), \end{aligned}$$

which are established as follows:

$$\begin{aligned} \sup h^*((-\infty, x)) &= \sup_{y < x} \sup h^{-1}((-\infty, y] \cap Z) \\ &= \sup h^{-1}(\overline{(-\infty, x)} \cap Z) \\ &= \sup h^{-1}((-\infty, x] \cap Z) \\ &= h^*(x) \\ \inf h^*((x, +\infty)) &= \inf_{y > x} \sup h^{-1}((-\infty, y] \cap Z) \\ &= \inf_{y > x} \inf h^{-1}((y, +\infty) \cap Z) \\ &= \inf h^{-1}((x, +\infty) \cap Z) \\ &= \sup h^{-1}((-\infty, x] \cap Z) \\ &= h^*(x). \quad \blacksquare \end{aligned}$$

Theorem 6. Let  $E$  be any subset of  $X$  and let  $\varphi: E \rightarrow R^q$  be any function of honorary Baire class  $2(E, R^q)$ . Then there exists a continuous function  $f: H \rightarrow R^q$  such that  $\varphi$  is a boundary function for  $f$ .



Proof. Let  $\psi : E \rightarrow \mathbb{R}^q$  be a function of Baire class 1 ( $E, \mathbb{R}^q$ ) and  $N$  a countable subset of  $E$  such that  $\varphi(x) = \psi(x)$  for every  $x \in E - N$ . Let  $\{s_n\}_{n=1}^{\infty}$  (with  $n \neq m$  implying  $s_n \neq s_m$ ) be a countable dense subset of  $X$  that includes every integer and every point of  $N$ . Let

$$t_n = \begin{cases} 1 & \text{if } s_n \text{ is an integer} \\ \frac{1}{2^n} & \text{if } s_n \text{ is not an integer.} \end{cases}$$

Define

$$h(x) = \begin{cases} \sum_{0 \leq s_n < x} t_n & \text{if } x > 0 \\ -\sum_{x \leq s_n < 0} t_n & \text{if } x \leq 0. \end{cases}$$

Then  $h$  is a strictly increasing function from  $\mathbb{R}$  into  $\mathbb{R}$ , and  $h(\mathbb{R})$  is bounded neither above nor below. Let  $h^*$  be the function described in Lemma 17.

Suppose that  $0 < y < 1$ . Then (for fixed  $x$ )

$$u - h^* \left( \frac{x - (1-y)u}{y} \right)$$

is a strictly increasing continuous function of  $u$  that approaches  $+\infty$  as  $u \rightarrow +\infty$  and  $-\infty$  as  $u \rightarrow -\infty$ . Consequently there exists precisely one number  $u(x, y)$  that satisfies the equation

$$(29) \quad u(x, y) - h^* \left( \frac{x - (1-y)u(x, y)}{y} \right) = 0.$$

I claim that  $u(x, y)$  is a continuous function on

$H_1 = \{ \langle x, y \rangle : x, y \in \mathbb{R} \text{ and } 0 < y < 1 \}$ . Suppose  $\{ \langle x_n, y_n \rangle \} \subseteq H_1$  and  $\langle x_n, y_n \rangle \rightarrow \langle x, y \rangle \in H_1$ . If  $u(x_n, y_n) \rightarrow +\infty$ , then

$$\frac{x_n - (1-y_n)u(x_n, y_n)}{y_n} \rightarrow_n -\infty,$$

and hence

$$u(x_n, y_n) - h^* \left( \frac{x_n - (1-y_n)u(x_n, y_n)}{y_n} \right) \rightarrow_n +\infty,$$

which contradicts (29). Thus  $u(x_n, y_n)$  cannot approach  $+\infty$ . A similar argument shows that  $u(x_n, y_n)$  cannot approach  $-\infty$ . Now assume that  $u(x_n, y_n) \rightarrow u_0 \in \mathbb{R}$ . Then, by (29),

$$\begin{aligned} 0 &= \lim_{n \rightarrow \infty} [u(x_n, y_n) - h^* \left( \frac{x_n - (1-y_n)u(x_n, y_n)}{y_n} \right)] \\ &= u_0 - h^* \left( \frac{x - (1-y)u_0}{y} \right), \end{aligned}$$

so  $u_0 = u(x, y)$ . By Lemma 16,  $u$  is continuous.

From Lemma 6, there exists a sequence  $\{g_n\}_{n=1}^{\infty}$  of continuous functions mapping  $X$  into  $\mathbb{R}^q$  such that  $g_n(x) \rightarrow \psi(x)$  for each  $x \in E$ . For  $n \geq 2$ , define

$$\begin{aligned} f_0(x, y) &= (yn(n+1) - n)g_n(u(x, y)) + ((n+1) - yn(n+1))g_{n+1}(u(x, y)) \\ &\quad \text{when } \frac{1}{n+1} \leq y \leq \frac{1}{n}. \end{aligned}$$

Then  $f_0$  is continuous on  $\bar{H}_2 \cap H$ . By the Tietze extension theorem, we can assume that  $f_0$  is defined and continuous on all of  $H$ . Let

$$\begin{aligned} r_n &= \inf_{x > s_n} h(x) \\ \ell_n &= \sup_{x < s_n} h(x) \\ v_n &= \varphi(s_n) - \psi(s_n) \quad \text{if } s_n \in N \\ v_n &= 0 \quad \text{if } s_n \notin N. \end{aligned}$$

If  $x$  and  $y$  are real numbers, define  $x \vee y = \max\{x, y\}$ . For  $\langle x, y \rangle \in H$ , set

$$\begin{aligned} \Delta_n(x, y) &= \\ &[(1 - ny) \vee 0] \left[ \left( 1 - \frac{1}{r_n - \ell_n} \left| r_n + \ell_n - 2s_n + 2 \frac{s_n - x}{y} \right| \right) \vee 0 \right] v_n. \end{aligned}$$

Then  $\Delta_n$  is continuous in  $H$ . Observe that  $\Delta_n(x, y) = 0$  when  $y \geq \frac{1}{n}$ .

Using this fact, it is easy to show that, if we set

$$f = f_0 + \sum_{n=1}^{\infty} \Delta_n,$$

then  $f$  is defined and continuous in  $H$ . We now show that  $\varphi$  is a boundary function for  $f$ .

Let  $p$  be any point of  $E$ . The line

$$(30) \quad x = (h(p) - p)y + p$$

passes through  $(p, 0)$ , and the part of it that lies in  $\bar{H}_1$  is an arc at  $p$ . We will show that  $f$  approaches  $\psi(p)$  along this line. If we substitute  $(h(p) - p)y + p$  for  $x$  in the expression for  $\Delta_n(x, y)$ , we obtain

$$(31) \quad \Delta_n(x, y) = [(1 - ny) \vee 0] \left[ \left(1 - \frac{1}{r_n - \ell_n} \left| r_n + \ell_n + 2\left(\frac{1}{y} - 1\right)(s_n - p) - 2h(p) \right| \right) \vee 0 \right] v_n.$$

If  $p \leq s_n$ , then  $h(p) \leq \ell_n$ , and one can verify directly that (31) vanishes. If  $p > s_n$ , then  $h(p) \geq r_n$ , and again one can verify directly that (31) vanishes. Thus  $\Delta_n(x, y)$  vanishes along that part of the line (30) lying in  $H$ .

Solving (30) for  $h(p)$ , we find that, along the given line,

$$h(p) = \frac{x - (1-y)p}{y},$$

and hence  $p = h^*(h(p)) = h^*\left(\frac{x - (1-y)p}{y}\right)$ . Therefore, if  $0 < y < 1$ ,  $p = u(x, y)$ . So, if  $\langle x, y \rangle$  satisfies (30),  $n \geq 2$ , and  $\frac{1}{n+1} \leq y \leq \frac{1}{n}$ , then

$$f_0(x, y) = (yn(n+1) - n)g_n(p) + ((n+1) - yn(n+1))g_{n+1}(p).$$

Since the coefficients of  $g_n(p)$  and  $g_{n+1}(p)$  in the above expression add up to 1 and since both coefficients lie in  $[0, 1]$ ,  $f_0(x, y)$  lies on the line segment joining  $g_n(p)$  to  $g_{n+1}(p)$ , and it follows that  $f_0(x, y)$  approaches  $\psi(p)$  as  $\langle x, y \rangle$  approaches  $p$  along the line (30). Since each  $\Delta_n$  vanishes on the part of this line lying in  $H$ ,  $f(x, y)$  also approaches  $\psi(p)$  along the line.

Let  $s_m$  be any point of  $N$ . We show that  $f$  approaches  $\varphi(s_m)$  along the part of the line

$$(32) \quad x = \left( \frac{r_m + \ell_m}{2} - s_m \right) y + s_m$$

that lies in  $H$ . Again, we first consider the value of  $\Delta_n$  along the given line. Substituting the value of  $x$  given by (32) into the expression for  $\Delta_n$ , we obtain

$$(33) \quad \Delta_n(x, y) = [(1-ny) \vee 0] \left[ \left( 1 - \frac{1}{r_n - \ell_n} |r_n - r_m + \ell_n - \ell_m + 2\left(\frac{1}{y} - 1\right)(s_n - s_m)| \right) \vee 0 \right] v_n.$$

If  $s_m < s_n$ , then  $\ell_m < r_m \leq \ell_n < r_n$ , and one can verify directly that (33) vanishes. If  $s_n < s_m$ , then  $\ell_n < r_n \leq \ell_m < r_m$ , and again one can verify that (33) vanishes. Thus, for  $n \neq m$ ,  $\Delta_n(x, y) = 0$  when  $\langle x, y \rangle$  lies on the line (32) and in  $H$ .

If we take  $n = m$  in (33), we obtain

$$\Delta_m(x, y) = [(1-my) \vee 0] v_m.$$

Therefore  $\Delta_m(x, y)$  approaches  $v_m = \varphi(s_m) - \psi(s_m)$  along the given line.

Take any  $\langle x, y \rangle \in H_1$  satisfying (32), and take any  $a$  and  $b$  satisfying

$$(34) \quad a < s_m < b.$$

Then  $h(a) \leq \ell_m < \frac{r_m + \ell_m}{2} < r_m \leq h(b)$ , so that

$(h(a) - s_m)y + s_m < x < (h(b) - s_m)y + s_m$ ; from which we deduce that

$$h(a) < \frac{x - (1-y)s_m}{y} < h(b).$$

Since  $h^*$  is weakly increasing,

$$a = h^*(h(a)) \leq h^*\left(\frac{x - (1-y)s_m}{y}\right) \leq h^*(h(b)) = b.$$

Since  $a$  and  $b$  were taken to be any two numbers satisfying (34), we conclude that

$$s_m = h^*\left(\frac{x - (1-y)s_m}{y}\right),$$

whence it follows that  $u(x, y) = s_m$ . Thus

$$f_0(x, y) = (yn(n+1) - n)g_n(s_m) + ((n+1) - yn(n+1))g_{n+1}(s_m)$$

when  $\langle x, y \rangle$  lies on the given line and  $\frac{1}{n+1} \leq y \leq \frac{1}{n}$ . Consequently

$f_0(x, y)$  approaches  $\psi(s_m)$  along the line (32). So  $f$  approaches  $\psi(s_m) + \varphi(s_m) - \psi(s_m) = \varphi(s_m)$ , and the theorem is proved. ■

**Theorem 7.** Let  $E$  be any subset of  $X$  and let  $\varphi: E \rightarrow S^2$  be any function of honorary Baire class 2( $E, S^2$ ). Then there exists a continuous function  $f: H \rightarrow S^2$  such that  $\varphi$  is a boundary function for  $f$ .

**Proof.** The proof of this theorem is very similar to that of Theorem 1.

Since  $S^2 \subseteq R^3$ , there exists, by Theorem 6, a continuous function

$g: H \rightarrow R^3$  having  $\varphi$  as a boundary function. Let

$$K = g^{-1}(\{v \in R^3 : |v| = \frac{1}{2}\})$$

$$L = g^{-1}(\{v \in R^3 : |v| \geq \frac{1}{2}\})$$

$$F = g^{-1}(\{v \in R^3 : |v| \leq \frac{1}{2}\}).$$

Let  $g_0 = g|_K$ .  $H$  is homeomorphic to  $R^2$ , so by [5, Lemma 2.9, p. 299],  $g_0$  can be extended to a continuous function

$$g_1 : H \rightarrow \{v \in R^3 : |v| = \frac{1}{2}\}.$$

Define  $f_1 : H \rightarrow R^3 - \{0\}$  by setting

$$f_1(z) = g(z) \quad \text{if } z \in L$$

$$f_1(z) = g_1(z) \quad \text{if } z \in F.$$

Then, since  $F$  and  $L$  are closed,  $f_1$  is continuous on  $H$ . It is easy to verify that  $\varphi$  is a boundary function for  $f_1$ . Let  $P_0 : R^3 - \{0\} \rightarrow S^2$  be the 0-projection onto  $S^2$  (see page 11), and let  $f$  be the composite function  $P_0 \circ f_1$ . Then  $f$  maps  $H$  continuously into  $S^2$ , and  $P_0 \circ \varphi = \varphi$  is a boundary function for  $f$ . ■

## CHAPTER II

### BOUNDARY FUNCTIONS FOR DISCONTINUOUS FUNCTIONS

#### 6. Boundary Functions for Baire Functions

It is not known whether the set of curvilinear convergence of a Borel-measurable function defined in  $H$  is necessarily a Borel set. The answer is not known even for functions of Baire class 1. However, a theorem on boundary functions that is similar to the corresponding result for continuous functions in  $H$  can be proved for functions of Baire class  $\xi$  in  $H$ .

**Definition.** If  $A$  and  $B$  are two sets, we will call  $A$  and  $B$  equivalent and write  $A \approx B$  if and only if  $A - B$  and  $B - A$  are both countable. It is easy to check that  $\approx$  is an equivalence relation.

**Lemma 18.** If  $A \approx E$ , then  $S - A \approx S - E$  for any set  $S$ . If  $A_n \approx E_n$  for all  $n$  in some countable set  $N$ , then

$$\bigcup_{n \in N} A_n \approx \bigcup_{n \in N} E_n \text{ and } \bigcap_{n \in N} A_n \approx \bigcap_{n \in N} E_n.$$

The proof of this lemma is routine.

**Definition.** An interval of real numbers will be called nondegenerate if it contains more than one point.

**Lemma 19.** Any union of nondegenerate intervals is equivalent to an open set.

Proof. Let  $\mathcal{I}$  be any family of nondegenerate intervals. It will suffice to prove that  $\bigcup_{I \in \mathcal{I}} I - \bigcup_{I \in \mathcal{I}} I^*$  is countable. We can write

$$\bigcup_{I \in \mathcal{I}} I^* = \bigcup_n J_n,$$

where  $\{J_n\}$  is a countable family of disjoint open intervals. If

$$x_0 \in \bigcup_{I \in \mathcal{I}} I - \bigcup_{I \in \mathcal{I}} I^*,$$

then  $x_0$  is an endpoint of  $I_0$  for some  $I_0 \in \mathcal{I}$ . For some  $n$ ,  $I_0^* \subseteq J_n$ ,

so that  $x_0 \in \bar{J}_n$ . But  $x_0 \notin J_n$ , so  $x_0$  is an endpoint of  $J_n$ . Thus

$\bigcup_{I \in \mathcal{I}} I - \bigcup_{I \in \mathcal{I}} I^*$  is contained in the set of all endpoints of the various  $J_n$ , and the lemma is proved. ■

Lemma 20. Let  $h$  be a weakly increasing real-valued function on a nonempty set  $E \subseteq \mathbb{R}$ . Suppose that  $|x - h(x)| \leq 1$  for every  $x \in E$ . Then  $h$  can be extended to a weakly increasing real-valued function  $h_1$  on  $\mathbb{R}$ .

Proof. Let  $e = \inf E$  ( $e$  may be  $-\infty$ ). For each  $x \in (e, +\infty)$ , set

$$h_1(x) = \sup h((-\infty, x] \cap E).$$

Since  $|t - h(t)| \leq 1$  for each  $t \in E$ ,

$$t \in (-\infty, x] \cap E \Rightarrow h(t) \leq x + 1,$$

so  $h_1$  is finite-valued. If  $e = -\infty$  we are done. If  $e > -\infty$ , then  $x \in E$  implies  $h(x) \geq x - 1 \geq e - 1$ , so  $h$  is bounded below. For  $x \in (-\infty, e]$  set

$$h_1(x) = \inf h(E).$$

It is easy to verify that  $h_1$  has the required properties. ■



Lemma 21. Let  $Y$  be a metric space,  $f : R \rightarrow Y$  a function of Baire class  $\xi(R, Y)$ , and suppose that  $h : R \rightarrow R$  is weakly increasing. Then there exists a countable set  $N \subseteq R$  such that the composite function  $f \circ h|_{R-N}$  is of Baire class  $\xi(R - N, Y)$ .

Proof. Let  $N$  be the set of discontinuities of  $h$ . By a well-known theorem,  $N$  must be countable. But then  $h|_{R-N}$  is continuous, so that  $f \circ (h|_{R-N}) = (f \circ h)|_{R-N}$  is of Baire class  $\xi(R - N, Y)$ . ■

Lemma 22. Let  $Y$  be a separable arcwise connected metric space,  $E$  any metric space, and let  $\varphi : E \rightarrow Y$  be a function having the following property. For every open set  $U \subseteq Y$  there exists a set  $T \in P^{\xi+1}(E)$  such that  $\varphi^{-1}(U) \subseteq T \subseteq \varphi^{-1}(\bar{U})$ . Then, if  $\xi \geq 2$ ,  $\varphi$  is of Baire class  $\xi(E, Y)$ .

Proof. The proof is similar to that of Lemma 15. Let  $\mathcal{B}$  be a countable base for  $Y$ , and suppose that  $W$  is any open subset of  $Y$ . Let

$$\mathcal{Q}(W) = \{U \in \mathcal{B} : \bar{U} \subseteq W\}.$$

The argument in the proof of Lemma 15 shows that

$$W = \bigcup_{U \in \mathcal{Q}(W)} U = \bigcup_{U \in \mathcal{Q}(W)} \bar{U}.$$

For each  $U \in \mathcal{B}$ , let  $T(U) \in P^{\xi+1}(E)$  be chosen so that  $\varphi^{-1}(U) \subseteq T(U) \subseteq \varphi^{-1}(\bar{U})$ . Then

$$\begin{aligned} \varphi^{-1}(W) &= \bigcup_{U \in \mathcal{Q}(W)} \varphi^{-1}(U) \subseteq \bigcup_{U \in \mathcal{Q}(W)} T(U) \\ &\subseteq \bigcup_{U \in \mathcal{Q}(W)} \varphi^{-1}(\bar{U}) = \varphi^{-1}(W). \end{aligned}$$

Thus  $\varphi^{-1}(W) = \bigcup_{U \in \mathcal{Q}(W)} T(U)$ , and since  $P^{\xi+1}(E)$  is closed under countable unions,  $\varphi^{-1}(W) \in P^{\xi+1}(E)$ . Therefore  $\varphi$  is of Baire class  $\xi(E, Y)$ . ■

Theorem 8. Let  $Y$  be a separable arcwise connected metric space,  $f : H \rightarrow Y$  a function of Baire class  $\xi(H, Y)$  where  $\xi \geq 1$ ,  $E$  a subset of  $X$ , and  $\varphi : E \rightarrow Y$  a boundary function for  $f$ . Then  $\varphi$  is of Baire class  $\xi + 1(E, Y)$ .

Proof. Let  $U$  be any open subset of  $Y$  and let  $V = Y - \bar{U}$ . Set

$$\begin{aligned} A &= \varphi^{-1}(U) & B &= \varphi^{-1}(V) \\ C &= A \cup B. \end{aligned}$$

Observe that  $A \cap B = \emptyset$ . For each  $x \in C$ , choose an arc  $\gamma_x$  at  $x$  such that

$$\begin{aligned} \lim_{\substack{z \rightarrow x \\ z \in \gamma_x}} f(z) &= \varphi(x) \\ \gamma_x &\subseteq \{z : |z - x| \leq 1\} \\ \gamma_x - \{x\} &\subseteq f^{-1}(U) & \text{if } x \in A \\ \gamma_x - \{x\} &\subseteq f^{-1}(V) & \text{if } x \in B. \end{aligned}$$

Notice that if  $x \in A$  and  $y \in B$ , then  $\gamma_x \cap \gamma_y = \emptyset$ .

We will say that  $\gamma_x$  meets  $\gamma_y$  in  $\bar{H}_n$  provided that  $\gamma_x$  and  $\gamma_y$  have subarcs  $\gamma_x'$  and  $\gamma_y'$  respectively such that  $x \in \gamma_x' \subseteq \bar{H}_n$ ,  $y \in \gamma_y' \subseteq \bar{H}_n$ , and  $\gamma_x' \cap \gamma_y' \neq \emptyset$ . Let

$$\begin{aligned} L_a &= \{x \in A : (\forall n)(\exists y)(y \in C, y \neq x, \text{ and } \gamma_y \text{ meets } \gamma_x \text{ in } \bar{H}_n)\} \\ L_b &= \{x \in B : (\forall n)(\exists y)(y \in C, y \neq x, \text{ and } \gamma_y \text{ meets } \gamma_x \text{ in } \bar{H}_n)\} \\ M_a &= \{x \in A : (\exists n)(\gamma_x \text{ meets no } \gamma_y \text{ (with } y \neq x) \text{ in } \bar{H}_n)\} \\ M_b &= \{x \in B : (\exists n)(\gamma_x \text{ meets no } \gamma_y \text{ (with } y \neq x) \text{ in } \bar{H}_n)\} \\ L &= L_a \cup L_b \\ M &= M_a \cup M_b. \end{aligned}$$

Observe that  $L_a, L_b, M_a, M_b$  are pairwise disjoint, and that  $A = L_a \cup M_a$  and  $B = L_b \cup M_b$ .

For each  $x \in M$ , let  $n(x)$  be a positive integer such that  $\gamma_x$  meets no  $\gamma_y$  (with  $y \neq x$ ) in  $\bar{H}_{n(x)}$ . Then  $n \geq n(x)$  implies that  $\gamma_x$  meets no  $\gamma_y$  in  $\bar{H}_n$ . Let

$$K_n = \{x \in C : \gamma_x \text{ meets } X_n, \text{ and, if } x \in M, n \geq n(x)\}.$$

Then  $K_n \subseteq K_{n+1}$  for each  $n$ , and  $C = \bigcup_{n=1}^{\infty} K_n$ .

We next show that for each positive integer  $n$  and each  $x \in L_a$  there exists a nondegenerate closed interval  $I_x^n$  such that  $x \in I_x^n \subseteq L_a \cup (X - K_n)$ . By the definition of  $L_a$ , there exists  $y \in C$  ( $y \neq x$ ) such that  $\gamma_y$  meets  $\gamma_x$  in  $\bar{H}_n$ . Let  $I_x^n$  be the closed interval having its endpoints at  $x$  and  $y$ . Let  $t$  be any point of  $I_x^n$ . We must prove that  $t \in L_a \cup (X - K_n)$ . If  $t \notin K_n$ , we are done. So assume  $t \in K_n$ . Then  $\gamma_t$  meets  $X_n$ , and hence it is clear from Figure 5 that  $\gamma_t$  must meet either  $\gamma_x$  or  $\gamma_y$  in  $\bar{H}_n$ . (This argument can be rigorized by means of Theorem 11.8 on p. 119 in [11].) But, if  $t \in M$ , then (because  $t \in K_n$ )  $n \geq n(t)$ , so that this situation is impossible. Therefore  $t \notin M$ . Now  $x \in L_a \subseteq A$ , so, since  $\gamma_x$  intersects  $\gamma_y$ ,  $y \notin B$ . Hence  $y \in C - B = A$ . Similarly, since  $\gamma_t$  intersects  $\gamma_x$  or  $\gamma_y$ ,  $t \in C - B = A$ . Thus  $t \in A - M = L_a$ , and we have shown that  $I_x^n \subseteq L_a \cup (X - K_n)$ .

Let  $W_n = \bigcup_{x \in L_a} I_x^n$ . For each  $n$ ,

$$L_a \subseteq W_n \cap C \subseteq [L_a \cup (X - K_n)] \cap C,$$

and therefore




Figure 5.

$$L_a \subseteq \left( \bigcap_{n=1}^{\infty} W_n \right) \cap C$$

$$\left\{ \bigcap_{n=1}^{\infty} [L_a \cup (X - K_n)] \right\} \cap C$$

$$= [L_a \cup (X - \bigcup_{n=1}^{\infty} K_n)] \cap C$$

$$= (L_a \cap C) \cup (C - \bigcup_{n=1}^{\infty} K_n) = L_a \cup \phi = L_a.$$

It follows that  $L_a = \left( \bigcap_{n=1}^{\infty} W_n \right) \cap C$ . By Lemma 19, each  $W_n$  is equivalent to an open set, so there exists a  $G_\delta$  set  $G_a \subseteq X$  such that

$$L_a \approx G_a \cap C.$$

A similar argument shows that there exists a  $G_\delta$  set  $G_b \subseteq X$  such that

$$L_b \approx G_b \cap C.$$

Next we study the properties of  $M_a$ . In doing this, it is convenient to define a function  $\pi : R^2 \rightarrow R$  by setting  $\pi(x, y) = x$ . If  $x \in M \cap K_n$ , then, starting at  $x$  and proceeding along  $\gamma_x$ , let  $p_n(x)$  be the first point of  $X_n$  that is reached. Define  $h_n^0 : M \cap K_n \rightarrow R$  by setting  $h_n^0(x) = \pi(p_n(x))$ . If  $x, x' \in M \cap K_n$  and  $x < x'$ , then, since  $\gamma_x$  cannot meet  $\gamma_{x'}$  in  $\bar{H}_n$ , it is evident that  $p_n(x)$  must lie to the left of  $p_n(x')$ ; that is,  $\pi(p_n(x)) < \pi(p_n(x'))$ . Thus  $h_n^0$  is a strictly increasing function on  $M \cap K_n$ . Moreover,

$$|x - h_n^0(x)| \leq 1 \quad \text{because } \gamma_x \subseteq \{z : |z - x| \leq 1\}.$$

So, by Lemma 20,  $h_n^0$  can be extended to a weakly increasing function  $h_n : X \rightarrow R$ . Let

$$g_n(x) = f(h_n(x), \frac{1}{n}) \quad (x \in R).$$

$f(x, \frac{1}{n})$  is a function (of  $x$ ) of Baire class  $\xi(X, Y)$ , so, by Lemma 21, there exists a countable set  $N_n \subset X$  such that  $g_n|_{X-N_n}$  is of Baire class  $\xi(X - N_n, Y)$ . Let  $N = \bigcup_{n=1}^{\infty} N_n$ . Then  $g_n|_{M-N}$  is of Baire class  $\xi(M - N, Y)$ .

For  $x \in M \cap K_n$ ,  $g_n(x) = f(h_n^0(x), \frac{1}{n}) = f(p_n(x))$ . If  $x \in M$ , then for all sufficiently large  $n$ ,  $x \in M \cap K_n$ , so

$$\lim_{n \rightarrow \infty} g_n(x) = \lim_{n \rightarrow \infty} f(p_n(x)) = \varphi(x).$$

Thus  $g_n|_M \rightarrow \varphi|_M$ , so  $g_n|_{M-N} \rightarrow \varphi|_{M-N}$ , hence  $\varphi|_{M-N}$  is of Baire class  $\xi + 1(M - N, Y)$ . It follows that there exists  $D \in P^{\xi+2}(X)$  such that

$$A \cap (M - N) = (\varphi|_{M-N})^{-1}(U) = D \cap (M - N).$$

Obviously  $A \cap M \approx D \cap M$ . Now,

$$L = L_a \cup L_b \approx (G_a \cap C) \cup (G_b \cap C) = (G_a \cup G_b) \cap C,$$

so

$$\begin{aligned} M_a &= A \cap M \approx D \cap M = D \cap (C - L) \\ &\approx D \cap [C - ((G_a \cup G_b) \cap C)] \\ &= D \cap [X - (G_a \cup G_b)] \cap C. \end{aligned}$$

$G_a$  and  $G_b$  are  $G_\delta$ , so  $X - (G_a \cup G_b)$  is  $F_\sigma$ , and hence

$$X - (G_a \cup G_b) \in P^2(X) \subseteq P^{\xi+2}(X).$$

Therefore  $M_a \approx F \cap C$ , where  $F \in P^{\xi+2}(X)$ . Now,  $G_a \in G_\delta(X) = Q^2(X)$ , and since  $\xi \geq 1$ ,  $Q^2(X) \subseteq P^{\xi+2}(X)$ , so  $G_a \cup F \in P^{\xi+2}(X)$ . But

$$A = L_a \cup M_a \approx (G_a \cap C) \cup (F \cap C) = (G_a \cup F) \cap C,$$

so  $A \approx S \cap C$ , where  $S \in P^{\xi+2}(X)$ . Since every countable set is  $F_\sigma$ , it is now easy to show that

$$A = T \cap C$$

for some  $T \in P^{\xi+2}(X)$ . From the definition of  $C$  it follows that

$T \subseteq X - B$ . Thus we have

$$\varphi^{-1}(U) = A \subseteq T \cap E \subseteq E - B = E - \varphi^{-1}(V) = \varphi^{-1}(\bar{U}).$$

$T \cap E \in P^{\xi+2}(E)$ , so Lemma 22 shows that  $\varphi$  is of Baire class  $\xi + 1(E, Y)$ . ■

Corollary. Let  $Y$  be a separable arcwise-connected metric space,

$f : H \rightarrow Y$  a Borel-measurable function,  $E$  a subset of  $X$ , and  $\varphi : E \rightarrow Y$  a boundary function for  $f$ . Then  $\varphi$  is Borel-measurable.

Proof.  $f$  is of some Baire class  $\xi(H, Y)$ , hence  $\varphi$  is of Baire class  $\xi + 1(E, Y)$ , hence  $\varphi$  is Borel-measurable. ■

This corollary raises the question of whether a boundary function for a Lebesgue-measurable function is necessarily Lebesgue-measurable, which we answer in the next section.

## 7. Boundary Functions for Lebesgue-Measurable Functions

Suppose that  $a_0, b_0, a_1, b_1$  are extended real numbers, and that  $a_0 \leq b_0, a_1 \leq b_1$ . To make the formalism more convenient we let  $(-\infty) - (-\infty) = 0$  and  $(+\infty) - (+\infty) = 0$ . In other respects we adhere to the usual conventions regarding arithmetic operations that involve  $-\infty$  or  $+\infty$ . Let

$$T(a_0, b_0; a_1, b_1) = \{ \langle x, y \rangle : 0 \leq y \leq 1 \text{ and } (a_1 - a_0)y + a_0 \leq x \leq (b_1 - b_0)y + b_0 \}.$$

A set of this form will be called a closed trapezoid. We also consider  $\phi$  to be a closed trapezoid. A set  $S$  will be called a trapezoid if there exists a closed trapezoid  $T$  such that  $T^1 \subseteq S \subseteq T$ ,

where  $T^1$  denotes the interior of  $T$  relative to  $\overline{H}_1$ . Every trapezoid is Lebesgue-measurable, though not necessarily Borel-measurable.

If  $s, s'$  are disjoint line segments having endpoints  $\langle a_0, 0 \rangle$ ,  $\langle a_1, 1 \rangle$ , and  $\langle a_0', 0 \rangle$ ,  $\langle a_1', 1 \rangle$  respectively, where  $a_i \leq a_i'$  ( $i = 0, 1$ ), then let

$$T(s, s') = T(s', s) = T(\overline{a_0, a_0'}; \overline{a_1, a_1'}).$$

If  $s = s'$ , then we let  $T(s, s') = T(s', s) = s$ . In what follows we will use the symbol  $X_0$  as an alternative designation for the  $x$ -axis  $X$ . This will enable us to make statements about  $X_i$  ( $i = 0, 1$ ) (where  $X_1$  denotes, as before,  $\{\langle x, 1 \rangle : x \in \mathbb{R}\}$ ).

We omit the proofs of the following two routine lemmas.

Lemma 23. Let the line segments  $s_1, s_2, s_3, s_4$  each have one endpoint on  $X_0$  and the other on  $X_1$ , and assume that  $i \neq j$  implies that either  $s_i \cap s_j = \emptyset$  or  $s_i = s_j$ . If  $T(s_1, s_2) \cap T(s_3, s_4) \neq \emptyset$ , then

$$T(s_1, s_3) \subseteq T(s_1, s_2) \cup T(s_3, s_4).$$

Lemma 24. Let  $\mathcal{S}$  be any set of line segments, each of which has one endpoint on  $X_0$  and the other on  $X_1$ , and no two of which intersect.

Then  $\bigcup_{s, s' \in \mathcal{S}} T(s, s')$  is a trapezoid.

Let  $m$  denote two-dimensional Lebesgue measure in  $\mathbb{R}^2$ . If  $E$  is a measurable subset of some line in  $\mathbb{R}^2$ , let  $m^k(E)$  denote the linear Lebesgue measure of  $E$ . Let  $\overline{m}_e$  and  $m_e^k$  denote two-dimensional exterior measure and linear exterior measure, respectively; i.e., for any  $E \subseteq \mathbb{R}^2$ ,

$$m_e(E) = \inf \{m(U) : E \subseteq U \text{ and } U \text{ is open}\};$$



and if  $E$  is a subset of a line  $L$ , then

$$m_e^{\mathcal{L}}(E) = \inf \{m^{\mathcal{L}}(U) : E \subseteq U \subseteq L \text{ and } U \text{ is open relative to } L\}.$$

Theorem 9. Let  $\mathcal{L}$  be any set of line segments, each of which has one endpoint on  $X_0$  and the other on  $X_1$ , and no two of which intersect.

Let  $S = \bigcup \mathcal{L}$ . Then

$$m_e(S) = \frac{1}{2} (m_e^{\mathcal{L}}(S \cap X_0) + m_e^{\mathcal{L}}(S \cap X_1)).$$

Proof. We may assume that  $\mathcal{L}$  is nonempty. Let  $\epsilon$  be any positive number. Choose an open set  $U \subseteq \mathbb{R}^2$  such that  $S \subseteq U$  and

$$m(U) \leq m_e(S) + \epsilon.$$

Let  $E_i = S \cap X_i$  ( $i = 0, 1$ ). Choose sets  $G_i \subseteq X_i$  that are open relative to  $X_i$  such that  $E_i \subseteq G_i$  and

$$m^{\mathcal{L}}(G_i) \leq m_e^{\mathcal{L}}(E_i) + \epsilon \quad (i = 0, 1).$$

Let  $V$  be the union of all lines  $L \subseteq \mathbb{R}^2$  such that  $L$  meets both  $G_0$  and  $G_1$ . It is easy to show that  $V$  is an open set. Furthermore,  $S \subseteq V$  and  $V \cap X_i = G_i$  ( $i = 0, 1$ ). Now let  $W = U \cap V$ . Then

$W$  is open,  $S \subseteq W \subseteq U$ , and

$$E_i \subseteq W \cap X_i \subseteq G_i \quad (i = 0, 1).$$

If  $s, s' \in \mathcal{L}$ , define  $s \equiv s'$  if and only if  $T(s, s') \subseteq W$ .

It is easy to verify by means of Lemma 23 that  $\equiv$  is an equivalence relation. Let  $\Gamma$  be the set of all equivalence classes. We prove that  $\Gamma$  is countable.

If  $s \in \mathcal{L}$ , we let  $\langle a_i(s), i \rangle$  be the endpoint of  $s$  on  $X_i$  ( $i = 0, 1$ ). Then

$$s = \{ \langle x, y \rangle \in \mathbb{R}^2 : 0 \leq y \leq 1 \text{ and } x = (a_1(s) - a_0(s))y + a_0(s) \}.$$

Since  $s$  is compact and contained in  $W$ , there is no difficulty in showing that there exists  $\delta_s > 0$  such that

$$\{ \langle x, y \rangle \in \mathbb{R}^2 : 0 \leq y \leq 1 \text{ and } \\ (a_1(s) - a_0(s))y + a_0(s) - \delta_s < x < (a_1(s) - a_0(s))y + a_0(s) + \delta_s \} \\ \subseteq W.$$

Let  $J_i(s) = (a_i(s) - \delta_s, a_i(s) + \delta_s)$  ( $i = 0, 1$ ). A sketch will rapidly convince the reader that if  $s, s' \in \mathcal{L}$ ,  $J_0(s) \cap J_0(s') \neq \phi$ , and  $J_1(s) \cap J_1(s') \neq \phi$ , then  $T(s, s') \subseteq W$ , so that  $s \equiv s'$ . Thus

$$(J_0(s) \times J_1(s)) \cap (J_0(s') \times J_1(s')) \neq \phi \Rightarrow s \equiv s'.$$

For each  $C \in \Gamma$ , choose  $s(C) \in C$  and let

$$Q(C) = J_0(s(C)) \times J_1(s(C)).$$

Then  $C_1 \neq C_2 \Rightarrow Q(C_1) \cap Q(C_2) = \phi$ . Since each  $Q(C)$  is a nonempty open subset of  $\mathbb{R}^2$ , this implies that  $\Gamma$  is countable.

If  $C \in \Gamma$ , let

$$T(C) = \bigcup_{s, s' \in C} T(s, s').$$

By Lemma 24,  $T(C)$  is a trapezoid. Also,

$$(35) \quad C \subseteq T(C) \subseteq W.$$

Suppose that  $C_1, C_2 \in \Gamma$  and  $C_1 \neq C_2$ . We claim that  $T(C_1) \cap T(C_2) = \phi$ . Assume that  $T(C_1) \cap T(C_2) \neq \phi$ . Then there exist  $s_1, s_1' \in C_1$  and  $s_2, s_2' \in C_2$  such that  $T(s_1, s_1') \cap T(s_2, s_2') \neq \phi$ . By Lemma 22,

$$T(s_1, s_2) \subseteq T(s_1, s_1') \cup T(s_2, s_2') \subseteq W,$$

so that  $s_1 \equiv s_2$ ; a contradiction. Therefore  $T(C_1) \cap T(C_2) = \phi$ .

Let  $K_i(C) = T(C) \cap X_i$  ( $i = 0, 1$ ). Then  $K_i(C)$  is an interval and

$$(36) \quad E_i \subseteq \bigcup_{C \in \Gamma} K_i(C) \subseteq W \cap X_i \subseteq G_i \quad (i = 0, 1).$$

Furthermore,  $C_1 \neq C_2$  implies that  $K_i(C_1) \cap K_i(C_2) = \phi$ . Using the formula for the area of a trapezoid, we find that

$$\begin{aligned} & \frac{1}{2} [m^k(\bigcup_{C \in \Gamma} K_0(C)) + m^k(\bigcup_{C \in \Gamma} K_1(C))] \\ &= \sum_{C \in \Gamma} \frac{1}{2} (m^k(K_0(C)) + m^k(K_1(C))) \\ &= \sum_{C \in \Gamma} m(T(C)) = m(\bigcup_{C \in \Gamma} T(C)). \end{aligned}$$

$$\begin{aligned} \text{Let } \alpha &= \frac{1}{2} [m^k(\bigcup_{C \in \Gamma} K_0(C)) + m^k(\bigcup_{C \in \Gamma} K_1(C))] \\ &= m(\bigcup_{C \in \Gamma} T(C)). \end{aligned}$$

According to (35),  $S \subseteq \bigcup_{C \in \Gamma} T(C) \subseteq W \subseteq U$ , so that

$$(37) \quad m_e(S) \leq \alpha \leq m(U) \leq m_e(S) + \epsilon.$$

By (36),

$$\begin{aligned} (38) \quad \frac{1}{2} (m_e^k(E_0) + m_e^k(E_1)) &\leq \alpha \leq \frac{1}{2} (m^k(G_0) + m^k(G_1)) \\ &\leq \frac{1}{2} (m_e^k(E_0) + m_e^k(E_1)) + \epsilon. \end{aligned}$$

Since  $\epsilon$  is arbitrary, inequalities (37) and (38) imply that

$$m_e(S) = \frac{1}{2} (m_e^k(E_0) + m_e^k(E_1)). \quad \blacksquare$$

One wonders to what extent a result resembling the foregoing theorem might be obtainable without the hypothesis that no two of the line segments intersect. The following example is relevant to this question. Let  $M_0$  be a residual set of measure zero in  $X_0$  and let  $M_1$

be a residual set of measure zero in  $X_1$ . Let  $\langle x_0, y_0 \rangle$  be any point of  $H_1$ . We claim that there is a line segment passing through  $\langle x_0, y_0 \rangle$  that has one endpoint in  $M_0$  and the other in  $M_1$ . For  $\theta \in (0, \pi)$ , let

$$\begin{aligned} F_1(\theta) &= \langle (1 - y_0) \cot \theta + x_0, 1 \rangle && \text{and} \\ F_0(\theta) &= \langle x_0 - y_0 \cot \theta, 0 \rangle. \end{aligned}$$

Then  $F_i$  is a homeomorphism of  $(0, \pi)$  onto  $X_i$ , so  $F_0^{-1}(M_0)$  and  $F_1^{-1}(M_1)$  are both residual sets in  $(0, \pi)$ . Choose  $\alpha \in F_0^{-1}(M_0) \cap F_1^{-1}(M_1)$ . Let  $L$  be the line whose equation is

$$x = x_0 + (y - y_0) \cot \alpha.$$

Then  $L$  passes through the points  $\langle x_0, y_0 \rangle$ ,  $F_0(\alpha)$  and  $F_1(\alpha)$ , so that  $L \cap H_1$  is the desired line segment. Let  $\mathcal{L}$  be the set of all line segments having one endpoint in  $M_0$  and the other in  $M_1$ . Let  $S = \bigcup \mathcal{L}$ .

Then  $S \cap X_0$  and  $S \cap X_1$  both have measure zero, but, as we have just shown,  $H_1 \subseteq S$ , so that  $S$  has infinite measure. See Problem 5 at the end of this paper.

Lemma 25. For every  $\epsilon > 0$  there exists a strictly increasing real-valued function  $h$  on  $\mathbb{R}$  such that  $h(\mathbb{R})$  has measure zero, and, for every real  $x$ ,  $|x - h(x)| \leq \epsilon$ .

Proof. For each integer  $n$ , let  $I_n = [n\epsilon, (n+1)\epsilon]$ . Then  $\bigcup_{n=-\infty}^{+\infty} I_n = \mathbb{R}$ . There exists a strictly increasing function  $f : [0, 1] \rightarrow [0, 1]$  such that  $m^k(f([0, 1])) = 0$ . For example, such a function may be defined as follows. Any number in  $[0, 1)$  may be written in the form

$$.a_1 a_2 a_3 \dots a_n \dots \quad (\text{binary decimal}),$$

where the decimal does not end in an infinite unbroken string of 1's.

Set

$$f(\cdot a_1 a_2 a_3 \dots a_n \dots) = \cdot b_1 b_2 b_3 \dots b_n \dots \quad (\text{ternary decimal}),$$

where  $b_i = 0$  if  $a_i = 0$  and  $b_i = 2$  if  $a_i = 1$ .

Set  $f(1) = 1$ . Then  $f$  maps  $[0, 1]$  into the Cantor ternary set, so  $m^k(f([0, 1])) = 0$ . It is easily shown that  $f$  is strictly increasing.

For each  $n$ , it is easy to obtain from  $f$  a function  $f_n : I_n \rightarrow I_n$  such that  $f_n$  is strictly increasing and  $m^k(f_n(I_n)) = 0$ . Set

$$h(x) = f_n(x) \quad \text{for } x \in (n\varepsilon, (n+1)\varepsilon].$$

There is no difficulty in proving that  $h$  has the required properties. ■

Theorem 10. There exists an indexed family  $\{\gamma_x\}_{x \in X}$  of simple arcs such that

- (i) for each  $x \in X$ ,  $\gamma_x$  is an arc at  $x$
- (ii)  $x \neq y \Rightarrow \gamma_x \cap \gamma_y = \emptyset$
- (iii)  $\bigcup_{x \in X} \gamma_x$  is a set of measure zero.

Proof. For each natural number  $n$ , let  $h_n : \mathbb{R} \rightarrow \mathbb{R}$  be a strictly increasing function such that  $h_n(\mathbb{R})$  has measure zero and, for every  $x$ ,  $|x - h_n(x)| \leq \frac{1}{n}$ . For every  $x \in \mathbb{R}$ , let  $s_n(x)$  be the line segment joining the point  $\langle h_n(x), \frac{1}{n} \rangle$  to the point  $\langle h_{n+1}(x), \frac{1}{n+1} \rangle$ . Since

$$h_n(x_1) < h_n(x_2) \Rightarrow x_1 < x_2 \Rightarrow h_{n+1}(x_1) < h_{n+1}(x_2),$$

we see that  $x_1 \neq x_2$  implies  $s_n(x_1) \cap s_n(x_2) = \emptyset$ . Let

$$S_n = \bigcup \{s_n(x) : x \in \mathbb{R}\}. \quad \text{Then}$$

$$S_n \cap X_n \subseteq \{ \langle x, \frac{1}{n} \rangle : x \in h_n(\mathbb{R}) \}$$

$$\text{and } S_n \cap X_{n+1} \subseteq \{ \langle x, \frac{1}{n+1} \rangle : x \in h_{n+1}(\mathbb{R}) \},$$

so  $m^k(S_n \cap X_n) = m^k(S_n \cap X_{n+1}) = 0$ . It is easy to deduce from

Theorem 9 that

$$m_e(S_n) = \left(\frac{1}{n} - \frac{1}{n+1}\right) \frac{1}{2} (m_e^l(S_n \cap X_n) + m_e^l(S_n \cap X_{n+1})) = 0.$$

For  $x \in X$ , let  $\gamma_x = \{x\} \cup \bigcup_{n=1}^{\infty} s_n(x)$ . Since  $\langle h_n(x), \frac{1}{n} \rangle \rightarrow x$ ,  $\gamma_x$  is an arc at  $x$ .

$$\begin{aligned} m_e\left(\bigcup_{x \in X} \gamma_x\right) &\leq m_e(X) + m_e\left(\bigcup_{n=1}^{\infty} S_n\right) \\ &\leq m_e(X) + \sum_{n=1}^{\infty} m_e(S_n) = 0, \end{aligned}$$

so  $\bigcup_{x \in X} \gamma_x$  is a set of measure zero. ■

Corollary. Let  $\varphi$  be an arbitrary function mapping  $X$  into any topological space  $Y$  having an element called 0. Then there exists a function  $f : H \rightarrow Y$  such that  $f(z) = 0$  almost everywhere and  $\varphi$  is a boundary function for  $f$ .

Proof. If  $\{\gamma_x\}_{x \in X}$  is the family of arcs described in Theorem 10, let

$$\begin{aligned} f(z) &= 0 && \text{if } z \text{ is in no } \gamma_x \\ f(z) &= \varphi(x) && \text{if } z \in \gamma_x. \end{aligned}$$

Then  $f$  is the desired function. ■

Corollary. There exists a real-valued Lebesgue-measurable function  $f$  defined in  $H$  having a nonmeasurable boundary function defined on  $X$ .

## SOME UNSOLVED PROBLEMS

1. If  $A$  is an arbitrary set of type  $F_{\sigma\delta}$  in  $X$ , does there necessarily exist a real-valued continuous function  $f$  defined in  $H$  having  $A$  as its set of curvilinear convergence? If  $\varphi$  is an arbitrary real-valued function of honorary Baire class 2 on  $A$  does there exist a continuous real-valued function  $f$  defined in  $H$  having  $A$  as its set of curvilinear convergence and  $\varphi$  as a boundary function?
  
2. (First proposed by J. E. McMillan [10]). If  $A$  is any set of type  $F_{\sigma\delta}$  in  $X$  and if  $\varphi$  is any function of honorary Baire class 2( $A, S^2$ ), does there necessarily exist a continuous function  $f : H \rightarrow S^2$  having  $A$  as its set of curvilinear convergence and  $\varphi$  as a boundary function?
  
3. If  $f$  is a real-valued Borel-measurable function defined in  $H$ , is the set of curvilinear convergence of  $f$  necessarily a Borel set? What if  $f$  is assumed to be of Baire class 1?
  
4. Let  $S = \{ \langle x, y, z \rangle \in R^3 : z > 0 \}$ . If  $f$  is a function defined in  $S$ , we define the set of curvilinear convergence of  $f$  in the obvious way. If  $f$  is continuous, is its set of curvilinear convergence necessarily a Borel set? Is it necessarily of type  $F_{\sigma\delta}$ ?
  
5. Let  $\mathcal{L}$  be a set of line segments each having one endpoint on  $X_0$  and the other on  $X_1$ , and let  $S = \bigcup \mathcal{L}$ . Assume that  $S$  is a Borel set. If  $m^k(S \cap X_0)$  and  $m^k(S \cap X_1)$  are known, what lower bound can be given

for  $m(S)$ ? A solution to this problem might be helpful in attacking a problem of Bagemihl, Piranian, and Young [3, Problem 1].



## REFERENCES

1. F. Bagemihl, Curvilinear cluster sets of arbitrary functions, Proc. Nat. Acad. Sci. U. S. A., 4 (1955) 379-382.
2. F. Bagemihl and G. Piranian, Boundary functions for functions defined in a disk, Michigan Math. J., 8 (1961) 201-207.
3. F. Bagemihl, G. Piranian, and G. S. Young, Intersections of cluster sets, Bul. Inst. Politehn. Iasi, 5 (9) (1959) 29-34.
4. S. Banach, Über analytisch darstellbare Operationen in abstrakten Räumen, Fund. Math., 17 (1931) 283-295.
5. S. Eilenberg and N. Steenrod, Foundations of algebraic topology, Princeton University Press, Princeton, New Jersey, 1952.
6. F. Hausdorff, Über halbstetige Funktionen und deren Verallgemeinerung, Math. Z., 5 (1919) 292-309.
7. F. Hausdorff, Set Theory, Second edition, Chelsea Publishing Company, New York, N. Y., 1962.
8. T. J. Kaczynski, Boundary functions for functions defined in a disk, J. Math. Mech., 14 (1965) 589-612.
9. T. J. Kaczynski, On a boundary property of continuous functions, Michigan Math. J., 13 (1966) 313-320.
10. J. E. McMillan, Boundary properties of functions continuous in a disc, Michigan Math. J., 13 (1966) 299-312.
11. M. H. A. Newman, Elements of the topology of plane sets of points, Cambridge University Press, 1964.
12. H. Tietze, Über stetige Kurven, Jordansche Kurvenbögen und geschlossene Jordansche Kurven, Math. Z., 5 (1919) 284-291.